

Left Radial Artery Access: A More Coaxial Approach to Cardiac Catheterization

An overview of our technique for left radial access, with a discussion of the benefits and limitations.

BY KEVIN FULLIN, MD, FACC, AND TIMOTHY SANBORN, MD, FACC, FAHA, FSCAI

In 2008, only 1% of cardiac catheterizations were performed using the radial approach, but this number grew to nearly 40% by 2017.¹ A preponderance of these radial approach cases were performed with a right radial approach because operators are familiar with working from the right side of the table and laboratory setup.² Several advantages of a left versus right radial approach have been documented, including reductions in contrast dose, fluoroscopy time, operator radiation exposure, needle-to-balloon time during acute myocardial infarction, and stroke.³⁻⁸ Another benefit of the

left radial approach is the ease of navigation through the radial/brachial/subclavian arteries, which can be done more coaxially as compared with the right radial approach.^{9,10} In addition, the left radial approach affords a more straightforward approach to left internal mammary artery and saphenous vein graft angiography.

For the left radial approach, some adjustments in technique are needed to compensate for ergonomic challenges for both the operator and patient. Difficulty of left-hand pronation has led to the distal radial approach. For instance, the anatomic snuffbox approach, pioneered by Dr. Ferdinand Kiemeneij, involves a catheter inserted distally into the hand between the patient's thumb and first finger. This has led to improved patient and operator comfort.¹¹ Other left radial artery techniques involve the left arm draped over the abdomen and supported by a Cobra board (TZ Medical Inc.).¹² This article describes an approach for a left radial technique and positioning for diagnostic catheterizations, coronary interventions, and rapid treatment of ST-segment elevation myocardial infarctions in which the patient's left arm is abducted out 90° from the body.

TECHNICAL DETAILS

The left radial artery access technique involves conventional placement of the patient on the catheterization laboratory table. The left arm is abducted 90° from the body and the hand is laid prone (Figure 1). A preparation table of similar height is positioned distal to the arm for preparing balloons, catheters, and stents. This approach emphasizes coaxial movement of catheters to the coronary



Figure 1. Left arm abducted to 90° allows coaxial catheter placement and substantial radiation protection of the operator's body and head. The first assistant is also removed from the radiation source with added radiation protection by use of the Zero-Gravity system.




Figure 2. The operator has excellent ergonomics during the procedure without back strain and torsion. The monitor screen is a short distance from the operator.



Figure 3. The technician who moves the table is remote from the radiation source and uses a radiation shield.

arteries and left ventricle. The operator stands on the caudal side of the extended left arm. The fluoroscopic and hemodynamic monitors are in front of the operator so that he or she is standing directly in front of the screens. There is no torsion on the operator's body, and he or she is looking forward at the monitors (Figure 2).

A catheterization technologist stands on the patient's right side and controls the table during the procedure using an additional monitor. A scrub technician stands to the right of the operator and prepares the equipment. One nurse circulates. An additional technician monitors hemodynamics and enters the patient's chronologic data.

The operator stands behind 800 lb of lead from the Zero-Gravity suspended radiation protection system (Biotronik, Inc.), which also protects the operator's head from radiation. The catheterization technician assistant also benefits from the Zero-Gravity radiation protection, which is positioned to the left of him or her (Figure 3). Real-time radiation sensors (RaySafe; Unfors RaySafe AB, a Fluke Biomedical company) are employed throughout the case. Access and closure with use of the Vasc Band hemostat (Teleflex) at the access site are accomplished without repositioning the arm.

LIMITATIONS

One limitation of the left radial approach is the requirement of four assistants, which is a manpower limitation in some catheterization laboratories. Alternatively, the left radial extended arm approach can be performed with three technicians if the x-ray controls are positioned on the left side of the table. Additionally, some creative solutions are necessary to have the manifold or assist device reach the extended left arm.

Using the Zero-Gravity system for radiation protection has some limitations in downward vision and the ability of

staff to clearly hear the operator's verbal directions. This can be solved by utilizing headphones with operative staff. If the Zero-Gravity system is not available, conventional aprons and vests can be used. This approach is less cumbersome but does not afford the same degree of radiation protection, particularly for the brain. The use of the Zero-Gravity system also affords its users less orthopedic stress because the protection is not worn but stepped into.

Additionally, the transradial approach may not be the best approach in every cohort of patients. A retrospective study showed decreased patency in radial arteries used for coronary artery bypass grafting that were previously utilized for angiography, as compared with those that had not been utilized during angiography.¹³ Evidence exists that the radial artery approach has an impact on vasomotor function, medial layer injury as documented by optical coherence imaging, and radial artery diameters postprocedure.¹⁴⁻¹⁶ Also, when using the radial artery for catheterization in patients with advanced kidney disease, the treating physician needs to consider the near-future use of that artery for possible creation of an arteriovenous dialysis fistula.¹⁷

CONCLUSION

This article describes a more coaxial way to perform heart catheterization and percutaneous interventions. This approach is supported by the literature, with reports of the left radial approach having fewer complications when compared with a right radial approach.¹⁸⁻²² The left-sided approach also enhances patient and operator comfort. We are unaware of previous reports of the left radial extended arm approach; however, it has been Dr. Fullin's standard approach for > 4 years. ■

(Continued on page 37)

(Continued from page 31)

1. Rao SV, Dharma S. 25 years of transradial intervention: looking back and anticipating what is ahead. *JACC Cardiovasc Interv.* 2017;10:2266-2268.
2. Rao SV, Cohen MG, Kandzari DE, et al. The transradial approach to percutaneous coronary intervention: historical perspective, current concepts, and future directions. *J Am Coll Cardiol.* 2010;55:2187-2195.
3. Xia SL, Zhang XB, Zhou JS, Gao X. Comparative efficacy and safety of the left versus right radial approach for percutaneous coronary procedures: a meta-analysis including 6870 patients. *Braz J Med Biol Res.* 2015;48:743-750.
4. De Rosa S, Torella D, Caiazzo G, et al. Left radial access for percutaneous coronary procedures: from neglected to performer? A meta-analysis of 14 studies including 7,603 procedures. *Int J Cardiol.* 2014;171:66-72.
5. Sciahbasi A, Romagnoli E, Burzotta F, et al. Transradial approach (left vs right) and procedural times during percutaneous coronary procedures: TALENT study. *Am Heart J.* 2011;161:172-179.
6. Fu Q, Hu H, Wang D, et al. Randomized comparative study of left versus right radial approach in the setting of primary percutaneous coronary intervention for ST-elevation myocardial infarction. *Clin Interv Aging.* 2015;10:1003-1008.
7. Rashid M, Lawson C, Potts J, et al. Incidence, determinants, and outcomes of left and right radial access use in patients undergoing percutaneous coronary intervention in the United Kingdom: a national perspective using the BCIS dataset. *JACC Cardiovasc Interv.* 2018;11:1021-1033.
8. Dominici M, Diletti R, Milici C, et al. Operator exposure to x-ray in left and right radial access during percutaneous coronary procedures: OPERA randomised study. *Heart.* 2013;99:480-484.
9. Kado H, Patel AM, Suryadevara S, et al. Operator radiation exposure and physical discomfort during a right versus left radial approach for coronary interventions: a randomized evaluation. *JACC Cardiovasc Interv.* 2014;7:810-816.
10. Norgaz T, Gorgulu S, Dagdelen S. A randomized study comparing the effectiveness of right and left radial approach for coronary angiography. *Catheter Cardiovasc Interv.* 2012;80:260-264.
11. Kiemeny F. Left distal transradial access in the anatomical snuffbox for coronary angiography (IdTRA) and interventions (IdTRI). *EuroIntervention.* 2017;13:851-857.
12. Casazza R. Distal left radial approach for cardiac catheterization: advancement in ergonomics with techniques and technologies. *Cath Lab Digest.* 2019;27:26-30.
13. Ruzieh M, Moza A, Siddagowda Bangalore B, et al. Effect of transradial catheterisation on patency rates of radial arteries used as a conduit for coronary bypass. *Heart Lung Circ.* 2017;26:296-300.
14. Mounsey CA, Mawhinney JA, Werner RS, Taggart DP. Does previous transradial catheterization preclude use of the radial artery as a conduit in coronary bypass surgery? *Circulation.* 2016;134:681-688.
15. Bhat T, Teli S, Bhat H, et al. Access-site complications and their management during transradial cardiac catheterization. *Expert Rev Cardiovasc Ther.* 2012;10:627-634.
16. Heiss C, Balzer J, Hauffe T, et al. Vascular dysfunction of brachial artery after transradial access for coronary catheterization: impact of smoking and catheter changes. *JACC Cardiovasc Interv.* 2009;2:1067-1073.
17. Hossain MA, Quinlan A, Heck-Kanellidis J, et al. Prevalence of chronic kidney disease among patients undergoing transradial percutaneous coronary interventions. *J Vasc Access.* 2018;19:346-349.
18. Kaledin A, Kochanov IN, Podmetin PS, et al. Distal radial artery in endovascular interventions. Research Gate website. https://www.researchgate.net/publication/319162208_Distal_radial_artery_in_endovascular_interventions. Published August 2017. Accessed June 24, 2019.
19. Shah B, Burdowski J, Guo Y, et al. Effect of left versus right radial artery approach for coronary angiography on radiation parameters in patients with predictors of transradial access failure. *Am J Cardiol.* 2016;118:477-481.
20. Dehghani P, Mohammad A, Bajaj R, et al. Mechanism and predictors of failed transradial approach for percutaneous coronary interventions. *JACC Cardiovasc Interv.* 2009;2:1057-1064.
21. Nishizaki Y, Yamagami S, Haga K, et al. Usefulness of prominently projected aortic arch on chest radiograph to predict severe tortuosity of the right subclavian or brachiocephalic artery in patients aged >44 years undergoing coronary angiography with a right radial artery approach. *Am J Cardiol.* 2012;110:203-207.
22. Cha KS, Kim MH, Kim HJ. Prevalence and clinical predictors of severe tortuosity of right subclavian artery in patients undergoing transradial coronary angiography. *Am J Cardiol.* 2003;92:1220-1222.

Kevin Fullin, MD, FACC

Froedtert South Medical Center

Kenosha, Wisconsin

kevin.fullin@froedtert-south.com

*Disclosures: None.***Timothy Sanborn, MD, FACC, FAHA, FSCAI**

Medical Director and Chair of Cardiology Services

Froedtert South Medical Center

Professor of Clinical Medicine

Medical College of Wisconsin

Kenosha, Wisconsin

tim.sanborn@froedtert-south.com

Disclosures: None.