


Endovascular TODAY

Fall 2015

VESSEL PREP IN THE NEW WORLD OF DRUG-ELUTION



Leading
endovascular
experts provide
their insights on
the evolving role
of atherectomy.



Robert
Beasley, MD



Jos C. van den
Berg, MD



Thomas P.
Davis, MD



Lawrence A.
Garcia, MD



William A.
Gray, MD



Akiko
Maehara, MD



J.A.
Mustapha, MD



Sonya S.
Noor, MD



Nicolas W.
Shammash, MD



Thomas M.
Shimshak, MD

Expert Panel Discussion on Atherectomy and Vessel Prep: What Roles Do They Play In My Practice?

Endovascular Today sat down with a multidisciplinary panel of esteemed interventionists to discuss their current practice paradigms for atherectomy, including the hot topic of Vessel Prep prior to drug-coated balloons and other adjunctive therapies.

MODERATOR



Craig McChesney

Publisher
Endovascular Today
Wayne, Pennsylvania
Mr. McChesney may be reached at
cmcchesney@bmctoday.com.

PARTICIPANTS



Robert Beasley, MS, MD

Director, Vascular/Interventional Radiology
MSMC Vein Center, and Wound Healing
Center at Mount Sinai Medical Center
Miami Beach, Florida

He has disclosed that he is a trainer and/or advisor for Abbott Vascular, Bard Peripheral Vascular, Boston Scientific Corporation, Cook Medical, Cordis Corporation, Cardiovascular Systems Inc, Endologix, Gore & Associates, Lake Region Medical, Lombard Medical, Medtronic, and Spectranetics Corporation.

Dr. Beasley may be reached at robert.beasley@msmc.com.



Thomas M. Shimshak, MD

Interventional Cardiologist
Heart and Vascular Center
Florida Hospital Heartland Medical Center
Sebring, Florida

He has disclosed that he receives honoraria from and is a consultant and speaker for Bayer, Volcano Corporation, and Boston Scientific Corporation, receives research support from Bayer, and does training programs for Bayer and Volcano Corporation.

Dr. Shimshak may be reached at tshimshak@gmail.com.



Sonya S. Noor, MD

Clinical Associate Professor
Department of Surgery
University of Buffalo
Buffalo, New York

She has disclosed that she is a consultant for Boston Scientific Corporation.

Dr. Noor may be reached at snoor@kaleidahealth.org.



Thomas P. Davis, MD

Director, Cardiac Catheterization Lab
St. John Hospital and Medical Center
Detroit, Michigan

He has disclosed that he is a consultant to Boston Scientific Corporation, Avinger, Roxwood Medical, Volcano, Medtronic, Cardiovascular Systems Inc., Cordis Corporation, and Bard Peripheral Vascular.

Dr. Davis may be reached at (313) 343-4612; tpdavis60@aol.com.



J.A. Mustapha, MD, FACC, FSCAI

Metro Health Hospital
Wyoming, Michigan
Michigan State University
Lansing, Michigan

He has disclosed that he receives honoraria from and is a consultant to Bard Peripheral Vascular, Boston Scientific Corporation, Cardiovascular Systems, Inc., Medtronic, and Spectranetics Corporation.

Dr. Mustapha may be reached at jihad.mustapha@metrogr.org.

All physicians were compensated for their time associated with this panel discussion and/or article.

EVT: In your practice, have you treated enough patients using the combination of atherectomy and drug-coated balloons (DCBs) (or other adjunctive therapies) to be able to say if your outcomes are better, worse, or about the same com-

pared to treatment with other contemporary therapies such as stenting, specialty balloons, etc.?

Dr. Mustapha: Luckily for us, we use ultrasound 100% in terms of bettering our therapy. One of the things that we've done so far [that] we've seen a difference with is debulk-

... for the first time, we have seen fewer patients coming back with restenosis since DCBs have been out, and atherectomy does make a big difference in those patients ...
—Dr. Mustapha

ing the target vessel pretty much to the wall, where you can actually go in, prep the vessel, deliver the DCB, and see under ultrasound that the DCB is actually in complete contact with the vessel wall, giving you a 1:1 ratio between the DCB and the vessel wall. I've got to tell you, for the first time, we have seen fewer patients coming back with restenosis since DCBs have been out, and atherectomy does make a big difference in those patients if you want to get a proper vessel prep prior to [using a] DCB.

Dr. Shimshak: I think the problem is, like all the trials, [what we see in the] real world is different in terms of the endpoints and the patient population that we deal with day in and day out. I think if we limit the scope of DCBs to short or intermediate-length lesions, there is clear benefit. What I struggle with is making the leap for long superficial femoral artery (SFA) disease—chronic total occlusions that are 20 or 30 cm. That is a huge amount of disease that you're dealing with. In those patients, I think atherectomy, coupled with whatever other device, gives you optimal luminal gain and is the way to go. That's why I think vessel preparation is so key. You may be able to achieve [good results in a long lesion] with a DCB, but I think that's a good place for stenting, and I still believe stenting has a role in those long segments of disease.

Dr. Davis: When we were in the DCB trials, we were treating a very different population of patients' lesions. Historically, I can say that I know a few patients who have come back when I've just used DCBs [to treat them]. I haven't seen many of the [patients treated with] atherectomy and DCBs, at least in my own sense of it, but we are collecting our data, and I think at the 1-year mark we'll know a little bit better. I think we need a little bit more time to go by to have a better answer to that question.

I think back over the European data and what Professor Thomas Zeller sees, and you've got to take what he says with a lot of stock. He truly is a believer in [atherectomy and DCB], and that's what he does in his practice.

Dr. Beasley: You see some tremendous results. Early on, I remember a couple of my patients who I treated with

atherectomy and DCBs when they first became available. Those patients have not shown any restenosis, reocclusion, or redevelopment of plaque that I can see, and these are patients that I follow closely on external ultrasound in the office.

In my lab, I'm using [multiple DCBs] at a time on a patient and hopefully getting a 70% to 80% success rate, where we don't have to place a stent. If we do have to place a stent, then you're placing a bare-metal stent over an area that already has drug on it.

EVT: What is the role of atherectomy in your current practice? Does it depend on the type of adjunctive therapy you are using?

Dr. Beasley: I use atherectomy in almost every case that has anything to do with any type of peripheral vascular disease—any type of critical ischemia or revascularization model. If [the vessel] has plaque, an occlusion, or a stenosis, I use atherectomy to prepare the vessel.

I'm a user of pretty much all the atherectomy devices, so depending on the location of the lesion, the position of the lesion, and the type of the lesion, I'll use a particular atherectomy device. I know with DCBs, you want to debulk and expose the vessel wall to that drug. With stenting, you want to give the stents a chance so that the stent's drug can appose itself to the wall the best possible way it can. So, I pretty much use atherectomy at all times.

Dr. Noor: I started using atherectomy early in my fellowship training so I took to it really easily—it wasn't as difficult to learn when you have already adopted other techniques, and at that time, it was really just angioplasty and stenting. I really like the philosophy of atherectomy, which has luminal gain and removal of the plaque, allowing the vessel to be more compliant with minimal trauma. Everything else that we do to the vessel in order to get luminal gain causes more injury and trauma and sets you back a year from now, when you have disease recurrence.

I use atherectomy, depending on the lesion, almost everywhere. In the femoral, popliteal, or below-the-knee distribution, atherectomy is probably my first line of defense. It then allows you to decide how you want to treat after. I'm not a big stenter; however, I will use focal stenting, depending on how much lesion or disease is left behind. I think it's a great platform for DCBs and possibly drug-eluting stents (DESs) once we have a little bit more data.

Dr. Mustapha: I try to marry each atherectomy device with the type of lesion or plaque that I'm facing at the time. In our institution, we use extravascular ultrasound, and that has been extremely helpful. We evaluate the plaque that we're dealing with and actually make a decision on which

type of atherectomy device we're going to use based on what we saw. [Under fluoroscopy,] we tend to undersize the vessel significantly, especially in the SFA/popliteal and tibial vessels. Based on what we see on ultrasound, we are able to debulk or modify the vessel [plaque] more accurately.

Dr. Shimshak: For me, it really comes down to plaque burden. When you begin to look at these vessels from a physical standpoint, you begin to realize why balloon angioplasty has failed at the most basic level. Remember, you cannot achieve an adequate lumen just by compressing that material. By the same token, even if you are an aggressive stenter, you [may] not be able to achieve optimal luminal dimensions of the stent even with the highest-performing stents today without preparing that vessel, in my opinion. I think balloon angioplasty is not enough. As Dr. Noor said, there is no question that as we begin to embark on preparing the vessel with balloon angioplasty when these vessels are highly calcified, there's a high incidence of dissections that are generated, which impacts patency and the durability of whatever intervention you perform.

Dr. Davis: To add onto that complexity, as years go on, we all push the envelope in terms of the types of vessels we're willing to treat. As the complexity of lesions increases, I

think atherectomy really is almost imperative in those types of lesions—long calcified lesions, thrombotic, and mixed lesions. As we push the limit, that's where we need it more.

Dr. Mustapha: Intimal calcification is different than medial calcification. Intimal calcification is the one that actually causes the problem for us, in particular if you deploy a stent without prepping the vessel properly—this is where you see the stent kinked. Intimal calcification has a significantly higher density of calcium deposit in it versus medial calcification. This is where the term “debulking” versus modifying the plaque comes into play. Knowing what kind of lesion you're dealing with is extremely important. Sometimes you may not be able to debulk it, so you modify it, and you will be able to dilate it.

Dr. Shimshak: To Dr. Mustapha's point, 360° calcification cannot be approached with angioplasty or any other device short of debulking. In my practice, the other traditional subset, as Dr. Davis alluded to, that we backed off from and now are pushing the envelope on, would be common femoral disease. I don't think every [case of] common femoral disease needs to be treated with endarterectomy. I think there are subsets that can be treated with endovascular techniques, and I think intravascular

ATHERECTOMY AND DCBs: A Q+A WITH DR. NICOLAS W. SHAMMAS



Nicolas W. Shammash, MD, MS, FACC, FSCAI

President and Research Director, Midwest Cardiovascular Research Foundation; Adjunct Clinical Associate Professor, University of Iowa; and Consultant and Interventional Cardiologist, Cardiovascular Medicine, PC Davenport, Iowa

He has disclosed that he received related research and educational grants for the Midwest Cardiovascular Research Foundation from Boston Scientific Corporation, Medtronic, and CSI; and is a trainer for the JETSTREAM Atherectomy System, is on the speakers' bureau for Boston Scientific Corporation, and is a member of the JET steering committee.

Dr. Shammash may be reached at shammash@mchsi.com.

EVT: In your practice, in which lesion types are you choosing to perform vessel prep with atherectomy prior to DCBs?

Dr. Shammash: I use atherectomy frequently prior to DCBs or plain old balloon angioplasty in any lesion > 10 cm long,

any total occlusions (if you cross intraluminally), or calcified plaque—irrespective of length.

Why do I do this? We have seen that these particular lesions are high risk for dissections and stenting, and I try to use a no-stent strategy in my lab as much as possible to keep the vessel intact for potential future treatments and avoid potential stent-related problems. With that no-stent strategy, atherectomy has become very important in my lab. If you look at the lesions that have the highest predictor for the need for stenting and the lesions that are more likely to dissect, they are calcified long lesions, total occlusions, and complex lesions (TASC C and D). With that in mind, these particular vessels are treated with atherectomy in my lab almost routinely. I have been performing atherectomy for over 15 years now, which has reduced my stenting rate to < 10%. Atherectomy is quite the tool to allow me to have the best acute procedural success.

EVT: Do you think thrombus is underappreciated in peripheral artery lesions? How important is clearing the thrombus barrier prior to utilizing DCBs?

Dr. Shammash: Clots can be of different ages, and different age clots bind paclitaxel in different ways and allow diffusion of the drug in different ways, creating a milieu that is highly unpredictable to how much drug can penetrate into the vessel

ultrasound (IVUS) is the guide for that, correlated with angiograms.

EVT: Despite a lack of level 1 data, what inspires your confidence in using atherectomy prior to DCB? What would you say to peers of yours who have not yet incorporated vessel prep with atherectomy into their practice, prior to using DCB in certain lesion types?

Dr. Noor: It's unfortunate that atherectomy doesn't have good level 1 evidence. It's probably a failure on all of our parts that there isn't good level 1 evidence that allows you to compare such a good modality of treatment with other standard modalities out there. But if I had to do every case with level 1 evidence, I would only get half my cases done, realistically.

It does take time. The problem with atherectomy is that there is a learning curve, and it's a steep learning curve. There are multiple devices out now, so it's difficult to be able to learn each one of them, but if you would pick one or two, you could use it. It's not as easy and fast as angioplasty and stenting—it's a labor of love—but I think you offer your patients a very good solution.

Dr. Beasley: When you take a look at IVUS and see the concentric luminal gain after atherectomy, and you take

wall. In my mind, the presence of a thrombus is equal to unpredictability of drug absorption into the vessel wall. If we can take care of the thrombus and remove it as much as possible, I think that would create more homogeneous, predictable drug diffusion into the target lesion.

With that in mind, I use the JETSTREAM™ Atherectomy System (Boston Scientific Corporation), because it's also approved for thrombectomy. I use it to treat both fibrous plaque and thrombus, and I try to remove as much of this plaque-thrombus burden as I can safely. The presence of a thrombus is also a high predictor for distal embolization, so particularly in total occlusions, I tend to use filters on a routine basis because we know very well that embolic debris will likely occur during the treatment.

EVT: How important is creating concentric lumens or circumferential lumens with atherectomy in order to create a uniform landing zone for DCB?

Dr. Shammass: My own particular preference is to maintain the rotational cutting within the intima and the superficial media rather than go deep into the media and the adventitia. There is a very interesting study that was recently published in the *Journal of Endovascular Therapy* that discussed the impact of deeper cuts into the media and the adventitia.¹ Strikingly, it showed that a very high rate of patency loss would occur when you cut very deep into the media and adventitia, supporting the hypothesis that restenosis, to a large degree, originates from

As the complexity of lesions increases, I think atherectomy really is almost imperative in those types of lesions—long calcified lesions, thrombotic, and mixed lesions. As we push the limit, that's where we need it more.

—Dr. Davis

the Fanelli results into account, and then the dispersion of the balloon and paclitaxel into the wall, when you improve the wall apposition—it's an argument you make without level 1 evidence, but it's an argument that I think that most reasonable folks can understand.

Dr. Shimshak: I think it's a leap at this point; we don't have the robust datasets that we want. I think it will come, but the message I would convey to people who are not yet embracing atherectomy, to help them understand the power of that therapy, would be to begin to use IVUS if they're not using IVUS. I think that is the key element in

the outer and deeper layers of the artery. To me, atherectomy is about vessel modification and about getting the least amount of deeper trauma into the vessel wall. I think this is a very important concept and raises the question of whether rotational cutting may lead to less restenosis than random directional cutting.

EVT: What type of clinical data would you like to see in the future regarding atherectomy and DCB?

Dr. Shammass: Peripheral vascular intervention lags behind the coronary world by years, and the reason for that is the lack of good, randomized data. I'd like to start seeing a move from just registries—which are also important—but, we need to move into the world of randomized trials that are powered enough to prove a point.

We need to be able to prove and get the message out there that atherectomy devices added to a DCB can be highly effective in reducing acute failure, and at the same time will likely have an impact on the long-term patency and reduction of target lesion revascularization. I would also like to see a trial of atherectomy with DCB versus DCB only that is powered and large enough to at least show that the additional vessel prep and the additional vessel modification would lead to better outcomes acutely as well as in the long term. ■

1. Tarricone A, Ali Z, Rajamanickam A, et al. Histopathological evidence of adventitial or medial injury is a strong predictor of restenosis during directional atherectomy for peripheral artery disease. *J Endovasc Ther.* 2015;22:712-715.

understanding the utility of this approach. Then, be guided by atherectomy coupled with other new technologies that do have more proven efficacy.

Dr. Mustapha: Stents did not always have level 1 evidence. Eventually, atherectomy will have level 1 evidence, and operators who don't use atherectomy today will hopefully see the value of atherectomy then. We had an atherectomy study [DEFINITIVE LE, Medtronic] that had a patency rate similar to stenting, so you already have something that tells you atherectomy is as effective as stenting in certain situations.

Many operators are reluctant to make the shift toward atherectomy utilization. In part, it could be due to the ease of use of a stent and/or not wanting to invest the time using atherectomy. In my opinion, atherectomy is the first tool to think of when trying to achieve the most effective vessel prep.

Dr. Davis: I go back to my Stone Age days when I was just using balloon angioplasty in the coronary [arteries], and then stents came out, and we always [thought] we shouldn't stent all the time, we should do bailout stenting. [Then] all of a sudden, stenting became this phenomenal thing. Then DESs came out, but because of the cost, you only used them in certain areas and at certain times. Now, if you don't put a DES in there, you're committing malpractice unless there's a good reason not to. So I think part of this is cost—costs have come down, and I think we've gotten used to the outcomes there.

I think drug delivery is here to stay, and right now, we have DCBs as our delivery system. Who knows where the technology is going to go and what's going to be the best delivery system, but that's what our system is.

EVT: How are you currently making the decision on which atherectomy device to use for vessel prep prior to using a DCB?

Dr. Shimshak: Calcium is critical to remove and prepare the vessel, but I also think it's plaque burden. Whatever device offers you the ability to debulk varying morphologies is my go-to device. There aren't very many that fit that bill. Most of the atherectomy devices are better for some things than others, but in my clinical practice, the JETSTREAM™ Atherectomy System (Boston Scientific Corporation) gives you predictability for varying lesion morphologies. Even for the non-IVUS users, I think there's comfort in that, if you don't understand the extent of disease, the device will perform admirably regardless of what kind of morphology you've encountered—soft plaque, eccentric, concentric, varying degrees of calcium, thrombus—it provides functionality for all those lesion morphologies.

... in my clinical practice, the
JETSTREAM Atherectomy System
gives you predictability for varying
lesion morphologies.

—Dr. Shimshak

In my practice, I would say over 90% [of the time, the] atherectomy device that I select off the shelf is JETSTREAM, for the reasons that I've already discussed. It gives me high performance for varying lesion morphologies, it's predictable, and it has a safety profile that's desirable. The aspiration is key, and I find it to be very desirable regardless of where I am.

Dr. Noor: Any time you performed a peripheral vascular intervention and you're concerned about thrombus, it's almost a contraindication to do anything because before, if you embolized the thrombus, we didn't have a lot of options. We had to lyse it and then you had to go back in and treat the underlying lesion.

But with newer technology, you can still go in and lyse it or use the AngioJet™ Thrombectomy System (Boston Scientific Corporation) and then treat it with atherectomy, or use JETSTREAM, which allows you to do both [atherectomy and thrombectomy, due to JETSTREAM having an indication for both]. The concept of being able to treat the thrombus and the underlying disease at the same time is very attractive not only for the patient and the time spent in the lab, but also from a cost standpoint. There are a lot of advantages to doing that, and your complication rate hopefully is lower with or without a filter, depending on how comfortable you are using one or not.

Dr. Davis: There is a lot more thrombotic disease that we just don't recognize in these lesions. Dr. Shimshak, as you've noticed, too, that's why by IVUS, with the virtual histology, you see it. When you do an OCT, the thrombotic areas are much more evident. A surgeon would probably recognize it more than we would, but as interventionists, you don't really recognize exactly how much thrombus you're dealing with on occlusive disease in the SFA.

Dr. Beasley: For the great majority of the SFA, [I use] rotational atherectomy because you have the benefit of not only being able to get a really nice channel, but also a very concentric luminal gain that you can then use for your adjunctive treatment modality, be it DCBs or stenting. You also have that aspirational component where you can at least be sure of yourself that you're pushing through and debulking this plaque. ■

The European Experience With Drug-Coated Balloons and Vessel Prep

With a 5-year head start using DCBs before US physicians, Prof. Jos van den Berg shares his take on best practices with this tool, as well as the available data.



Jos C. van den Berg, MD, PhD

Head of Service of Interventional Radiology
Ospedale Regionale di Lugano, cede Civico
Lugano, Switzerland

Associate Professor of Radiology

University of Bern, Switzerland

He has stated that he has no financial interests related to this article.

Prof. van den Berg may be reached at jos.vandenberg@eoc.ch.

EVT: As someone who has had the opportunity to use drug-coated balloons (DCBs) for a number of years, what would you say to the US physicians who are looking to understand how well DCBs work?

Prof. van den Berg: When we started using DCBs 5 years ago, we actually didn't know anything about how they worked or whether they worked. So in that respect, being a little bit late may be an advantage, because we know now how they work. When we started, not having any data, we were just using those balloons in difficult lesions. We now have, more or less, the picture that it also works in primary lesions—at least TASC A and B.

EVT: Is that knowledge you've gleaned from the randomized studies or from your own practice?

Prof. van den Berg: It's mainly from the randomized studies. Seeing that it worked in my own practice also helps, but I think when you have the confirmation from larger randomized trials, then you can really be confident that the new technique is working.

EVT: From a clinical standpoint, what are some of the salient points about vessel prep that you have learned over the past several years?

Prof. van den Berg: Vessel preparation can be very

helpful. In patients who don't have [a lot of] calcification in the vessel wall, you can do it with just optimal balloon angioplasty. In cases where patients have highly calcified SFAs, then you probably need to do something additional [such as atherectomy].

Regarding optimal balloon angioplasty, I think it's important to be very meticulous with your technique. A lot of people just inflate a balloon rapidly and deflate very rapidly, and that's one of the things that probably enhances the incidence of restenosis by creating this trauma to the vessel wall. By gently inflating the balloon, you really give the vessel some time to adapt to the balloon, not creating much vessel wall injury. By leaving the balloon inflated for a long time, recoil will probably be much less. We know this from studies in the past, in the 1980s and 1990s, when stents were not available, and people had balloon angioplasty as the only tool. Even in the long lesions with long dissections, you can actually get rid of the complication of dissection with a long balloon inflation. That is something we had forgotten about when long stents became available for the SFA.

EVT: In heavily calcified lesions, what are some of the additional steps that you try to take?

Prof. van den Berg: One of the problems with the heavily calcified lesions, as we know from the Fanelli study, is that the calcium is really interfering with the good results of DCBs.¹ You probably need to get rid of (or crack) the calcium in order to enhance the results in those specific patients. Fanelli et al made a classification from grade 1 to 4, and they really saw a drop off in primary patency in the grade 4 lesions that had calcium all around the vessel wall, almost at 360°. So, the idea is to get that out in order to get better results with the DCBs.

There are some data from a small study from Cioppa et al² in Italy that indicate that by using atherectomy to take out the calcium and then following up with a DCB

(Continued on page 22)

The Data Behind Atherectomy and Drug-Coated Balloons

Dr. Lawrence A. Garcia shares his thoughts on what is needed most from future trials and reflects on his key learnings in this space.



Lawrence A. Garcia, MD

Section of Interventional Cardiology and
Peripheral Interventions
St. Elizabeth's Medical Center
Boston, Massachusetts.

He has disclosed that he does research for and is a noncompensated consultant for Abbott Vascular and Medtronic; is a noncompensated consultant for Bard Peripheral Vascular, Cook Medical, TriVascular, and Boston Scientific Corporation; is a compensated consultant for St. Jude Medical; and has equity in CV Ingenuity, Primacea, Spirox, Tissue Gen, Arsenal, Syntervention, Essential Medical, and Scion Cardiovascular.

Dr. Garcia may be reached at lawrence.garcia@steward.org.

EVT: What type of trial data do you want to see moving forward in the atherectomy and drug-coated balloon (DCB) space, and why? What type of “real world” questions might the data help to answer?

Dr. Garcia: What we really need now is head-to-head comparisons. At the end of the day, everybody is going to be looking for superiority trials.

I suspect that for a superficial femoral artery (SFA) or an infrapopliteal segment, which is so inhospitable for everything we do endovascularly, that as long as you get a primary patency and the pattern of restenosis is less aggressive than what you first started with, then the therapy for that is a lot cheaper. What you spend up front becomes absolutely critical to the health care dollar downstream, and that's where I think we're going to win or lose on a lot of these head-to-head trials.

The combination therapy of atherectomy and DCB has a large cost, but if the downstream side is that you have a 91% primary patency, and of those 9% that fail, they fail in focal ways, then the downstream reintervention for any patient who needs it is a balloon. [This is more cost efficient] as opposed to having a similar up front cost [with a stent] and having a failure that may become an occlusion, which may then require more

expensive reintervention, so your health care dollar just got wasted. I think that is where we're going with ACO models and primary payers—we're going to have to focus a lot on the health care economics.

EVT: You have been intimately involved in both atherectomy and DCB trials. What have you learned in your experience that you're applying to your practice today about both of those?

Dr. Garcia: We've championed atherectomy for a lot of years, and I still believe that the technology, in and of itself, particularly for the SFA, is a very viable and valuable commodity for how we treat our patients when it comes to treatment for claudication. The DCB world, I think, is the holy grail. A lot of us in the United States have seen other parts of the world, particularly Europe and Asia, get DCBs for so many years, and we felt left out. However, once DCBs got here, I think many of us have gravitated to using them in these anatomic locations, particularly the SFA.

In my particular practice, I have found that the combination of therapy, both with atherectomy as well as with DCB is very useful. My hope is that our anecdotal experience translates scientifically when testing combination therapy versus DCBs alone, or against what should be considered the standard right now, which is Zilver PTX's (Cook Medical) 5-year data. Eventually, we have to go against other therapies, and if the endoprostheses wins, then it will save a lot of time, but we should prove it. If it fails, and atherectomy is proven to be best with the combination therapy, then we should gravitate toward that.

In my particular practice, I've always been somebody who likes to leave nothing behind, and it's interesting to see the worldwide consensus come back to the folks who used to stent a ton and now say that they are leaving nothing behind. I think we've all learned that once you put a stent in there, it's in there forever, and you have to deal with it in some way, shape, or form in the future. ■

Effective Debulking With the JETSTREAM™ Atherectomy System

Strategies for effective treatment of PAD in the era of drug-coated balloons and contemporary stenting.

BY THOMAS M. SHIMSHAK, MD, FACC, FSCAI

Peripheral artery disease (PAD) is a major cause of morbidity and mortality in the United States, affecting 8 to 12 million people. The incidence of PAD increases in the presence of well-defined atherosclerotic risk factors, including cigarette smoking, diabetes mellitus, hypertension, hyperlipidemia, and advanced age, and is estimated to affect > 20% of adults aged 55 years and older. When symptomatic, PAD may adversely have an impact on functional capacity, ability to work, and quality of life. Furthermore, PAD is associated with significant social and economic costs¹ and increases the risk of future cardiovascular events.

Advances in percutaneous catheter-based therapies have led to improved early and late clinical results in symptomatic patients.² Successful percutaneous revascularization improves quality-of-life measures, functional capacity, amputation rates, and survival in patients with intermittent claudication and critical limb ischemia. Use of adjunct devices and improved procedural outcomes have resulted in an increase in the number of PAD patients treated with endovascular therapy. The number of endovascular procedures has doubled for patients with intermittent claudication, and it has increased fourfold in patients with critical limb ischemia.²

Endovascular therapy of the superficial femoropopliteal arterial segment has historically been challenging. Although overall procedural results have been favorable, late results have been limited by unacceptable high restenosis rates and recurrent symptoms. The atherosclerotic disease process in the femoropopliteal arterial segment is often diffuse with complex histologic morphologies, including soft or fibrous tissue, thrombus, and superficial and deep calcium. In addition, chronic total occlusions (CTOs) are common (Table 1).³ These factors have limited the utility of balloon angioplasty alone for sustainable favorable results and have led to the use of alternative therapies, includ-

TABLE 1. LESION LOCATION AND CHARACTERISTICS*

Variable	No. (%) or Mean \pm SD
Stenosis	1,334 (62.4)
Chronic total occlusion	615 (28.8)
In-stent restenosis	188 (8.8)
Mean length, mm	100.8 \pm 9.4
Location	
Femoral	660 (30.9)
Popliteal	266 (12.4)
Tibial	513 (24.0)
Bypass graft	59 (2.7)
Multilevel	389 (18.3)
TASC classification	
A	297 (13.9)
B	632 (29.5)
C	592 (27.7)
D	616 (28.8)

*Based on data from Shrikhande GV, Khan SZ, Hussain HG, et al. Lesion types and device characteristics that protect distal embolization during percutaneous lower extremity interventions. *J Vasc Surg*. 2011;53:347-352.³

ing stenting (bare-metal, drug-eluting, and covered nitinol stents), atherectomy, and more recently, drug-coated balloons (DCBs).

The JETSTREAM Atherectomy System (Boston Scientific Corporation) is intended for use in atherectomy of the peripheral vasculature and to break apart and remove thrombus. It consists of a sterile, single-use catheter and control pod and a reusable power console. The catheter is compatible with an 0.014-inch wire (including the Thruway™ Guidewire [Boston Scientific




Figure 1. The JETSTREAM Atherectomy System cutters. Two different expandable cutter (XC) catheters are available in sizes 2.1/3.0 mm and 2.4/3.4 mm. The single cutter (SC) catheters have fixed, nonexpandable cutters and are available in two sizes, 1.6 mm and 1.85 mm. All catheters are 7-F sheath and 0.014-inch guidewire compatible (including the Boston Scientific Thruway Guidewire. It is also approved for use with Atherectomy Lubricants, such as Rotaglide™ Lubricant).

Corporation]) and 7-F sheath. It consists of a five-flute, front-end cutting tip that rotates at 70,000 to 73,000 RPM. Catheters are available in a variety of sizes, including two with nonexpandable cutters (1.6 mm or 1.85 mm), and two catheters with expandable cutters (2.1/3.0 mm, and 2.4/3.4 mm) (Figure 1). The JETSTREAM Atherectomy System is designed to treat a variety of lesion morphologies including soft, fibrotic, calcified, and/or thrombus. By virtue of its property of differential cutting, it preferentially cuts atheromatous disease, while sparing normal tissue. It also incorpo-

rates dynamic and continuous aspiration of particulate debris and thrombus, a feature that reduces distal emboli and improves device and procedural safety. Although other atherectomy systems have demonstrated effectiveness in removing calcium, the JETSTREAM Atherectomy System is unique in terms of combining differential cutting with dynamic aspiration (Table 2).

CASE PRESENTATIONS*

Case 1: Diffuse Distal SFA and Popliteal CTO

A 78-year-old woman presented with severe, limiting, intermittent claudication of her right leg. She had undergone complex endovascular therapy of her left leg several months earlier, after presenting with an ischemic great toe ulcer. The ulcer had healed, but she had limiting exertional right calf pain, which had been present for more than 6 months. Previous CT angiography had demonstrated wide patency of the right common and external iliac arteries, common femoral artery (CFA) and profunda, and proximal right superficial femoral artery (SFA). The distal right SFA was diffusely diseased, and the popliteal artery was chronically occluded.

The interventional procedure was completed using antegrade access with a 7-F sheath, demonstrating a diffusely diseased, calcified distal right SFA with multiple subtotal stenoses. The proximal portion of the popliteal artery was chronically occluded with reconstitution of the midportion of the popliteal artery via collaterals (Figure 2). The distal popliteal artery had a dis-

TABLE 2. COMPARISON OF PROPERTIES OF DIFFERENT ATHERECTOMY DEVICES

	JETSTREAM™ Atherectomy System (Boston Scientific Corporation)	Peripheral Rotablator™ Rotational Atherectomy System (Boston Scientific Corporation)	Diamondback 360™, Stealth 360™ Atherectomy System (Cardiovascular Systems, Inc.)	SilverHawk™, TurboHawk™ Plaque Excision System (Medtronic)	Turbo- Elite Laser™ Atherectomy Catheter (Spectranetics Corporation)
Front cutting	✓	✓			N/A
Differential cutting	✓	✓	✓		N/A
Active aspiration	✓				
Concentric lumens	✓	✓			
Lesion morphology:					
Calcium	✓	✓	✓	✓ (large vessel only)	✓
Thrombus	✓				✓
Sources: Endovascular Today Buyer's Guide 2014. JETSTREAM System Brochure, Boston Scientific Website, 2014. Peripheral Rotablator product website, Boston Scientific, 2014. Diamondback 360 product website, CSI, 2014. Covidien website, Directional Atherectomy products, 2014. Turbo-Elite Laser Atherectomy Catheter Instructions for Use, May 2014.					

*Results from case studies are not necessarily predictive of results in other cases. Results in other cases may vary.




Figure 2. Distal right SFA (A). CTO of the proximal popliteal artery (B). Reconstituted midpopliteal artery (C).




Figure 3. IVUS using virtual histology of the popliteal artery following recanalization and predilatation. Eccentric dense fibrocalcific plaque with scattered thrombus and an extensive arc of superficial calcium encompassing $> 270^\circ$ (A). Eccentric fibrocalcific plaque and scattered associated thrombus (B). Predominantly fibrous plaque and localized thrombus and superficial calcium (C).

crete subtotal stenosis. There was diffuse infrapopliteal disease, characterized by an occluded distal peroneal artery and proximal occlusion of the posterior tibial artery. The anterior tibial artery was widely patent.

After administering 2,500 units of intravenous (IV) heparin, a 0.018-inch, 300-cm-long, 30-g Victory™ Guidewire (Boston Scientific Corporation) was used to recanalize the CTO using a 4-F angled Glidewire® (Terumo Interventional Systems). The wire tip was directed freely into the anterior tibial artery with fluoroscopic guidance. After recanalizing the occluded popliteal artery, an additional 5,000 units of IV heparin were administered to achieve a therapeutic activated clotting time. After exchanging for a 0.014-inch guidewire, the popliteal artery and distal SFA were then dilated with a 2-X 150-mm balloon catheter, performing multiple overlapping inflations encompassing the mid and proximal popliteal artery and distal SFA. The 0.018-inch guidewire was exchanged for a 0.014-inch, 315-cm-long BareWire (Abbott Vascular) delivery wire. Intravascular ultrasound (IVUS) was then performed using the 2.5-mm Eagle Eye® Platinum IVUS catheter (Volcano Corporation). Virtual

histology and Chromaflo (Volcano Corporation) were used to assess the disease severity, extent of calcium, lesion morphology, and vessel dimensions and found severe fibrocalcific disease with mixed thrombus and extensive superficial calcium encompassing a $> 270^\circ$ arc of calcium (Figure 3).

The JETSTREAM Atherectomy System was used to debulk the lesion and remove calcium. Prior to introducing the 1.6-mm cutter, a 4- to 7-mm Emboshield NAV6® (Abbott Vascular) was deployed in the distal popliteal artery. The 1.6-mm cutter was advanced manually just proximal to the disease. The device was then activated, and two passes were made, encompassing the entire length of the diseased segment. This was followed by additional passes with the 2.1- to 3-mm cutter. Adjunctive angioplasty of the distal SFA and proximal popliteal arteries was performed with a 4-mm Chocolate® balloon (Cordis Corporation). A 4.5-mm braided self-expanding stent was deployed in the distal SFA with minimal elongation. Final angiogram and IVUS demonstrated wide patency and full stent apposition (Figures 4 and 5). The distal popliteal lesion was also




Figure 4. The JETSTREAM Atherectomy System was used on the right distal SFA, which had associated calcium (A). Final arteriogram of distal SFA after using the JETSTREAM Atherectomy System, angioplasty, and stenting (B). Final arteriogram of the popliteal artery after using the JETSTREAM Atherectomy System and balloon angioplasty (C).




Figure 5. IVUS images of the popliteal artery after JETSTREAM Atherectomy, adjunctive balloon angioplasty (A), and stenting (B). Posttreatment IVUS showed significant calcium and plaque removal and an increase in luminal cross-sectional area compared to pretreatment.




Figure 6. Angiogram of the right CFA, SFA, and profunda artery (A). IVUS images of the CFA (B) and SFA origin (C).

treated with the 2.1-mm JETSTREAM Atherectomy System. The patient's postprocedure course was uneventful. She was discharged the following day and remained stable and symptom free at 6-month follow-up.

Case 2: Complex Calcific Disease in the CFA, Profunda, and SFA

An 84-year-old man presented with severe, bilateral exertional calf discomfort after walking < 1 block. Noninvasive evaluation included a resting ankle-brachial index in the right leg of 0.68. He underwent CT angiography, which demonstrated discrete, high-grade disease of the right CFA and of the proximal and distal right SFA. He had two-vessel runoff below the knee, consisting of the anterior tibial and peroneal arteries. He was referred for selective right iliofemoral arteriography and runoff and possible endovascular therapy. Contralateral access from the left femoral artery using a 6-F Flexor® Ansel sheath (Cook Medical) was used. Selective right iliac arteriography demonstrated highly calcified, complex disease and critical obstructive disease of the right SFA with an eccentric subtotal (95%) stenosis. There was also high-grade disease at the origin of the profunda artery, with an eccentric 85% to 90% stenosis. The CFA was extensively calcified and had significant distal disease just proximal to the bifurcation (Figure 6). The patient was given 5,000 units of IV heparin, and the right SFA was crossed with a 0.014-inch 30-g Victory™ Guidewire. A second 0.014-inch, ChoICE™ PT Guidewire (Boston Scientific Corporation) was directed into the profunda artery using a two-wire technique. A series of sequential balloon inflations were then performed with 2.5-mm and 3-mm X 20-mm balloons, dilating the CFA-SFA origin, followed by the CFA-profunda arteries. The balloons were inflated to 10 to 12 atm, and full balloon expansion was achieved.

At this point, the CFA, SFA, and profunda artery were evaluated with the 2.5-mm Eagle Eye® Platinum IVUS catheter. IVUS demonstrated a dense, extensive arc of superficial calcium of the SFA, encompassing > 270° of the luminal circumference (Figure 6). Based on the arteriographic findings and the IVUS data, the JETSTREAM Atherectomy System was used. A 4-F Glidecath® was advanced over the 0.014-inch-long guidewire into the mid-SFA. The 0.014-inch guidewire was exchanged for a 0.035-inch, 300-cm Supra Core® (Abbott Vascular) guidewire. The 6-F sheath was then exchanged for a 7-F Flexor Ansel® contralateral sheath, which was advanced to the distal right external iliac artery. An additional 2,500 units of IV heparin were administered to achieve a therapeutic activated clotting time. The Glidecath® was reintroduced over the Supra Core® wire to the




Figure 7. The 1.85-mm SC JETSTREAM cutter (A) and 2.1- to 3-mm XC JETSTREAM cutter (B). Balloon angioplasty of the proximal CFA-SFA after JETSTREAM Atherectomy (C). Balloon angioplasty of the CFA-profunda artery after JETSTREAM Atherectomy (D).

mid-SFA. The 0.035-inch wire was then exchanged for a 0.014-inch, 315-cm BareWire, which was advanced to the distal SFA, and a 4- to 7-mm Emboshield NAV6[®] was then deployed. The guidewire in the profunda artery was removed. The CFA and origin of the SFA were then treated with the 1.85-mm SC JETSTREAM catheter, followed by the 2.1- to 3-mm XC JETSTREAM catheter (Figure 7).

After withdrawing the JETSTREAM catheter over the wire, repeat arteriography demonstrated significant improvement in the SFA, with < 50% residual disease. There was persistent high-grade disease at the origin of the profunda artery. The SFA was then dilated with a series of prolonged, low-pressure inflations using the 5-mm balloon (Figure 7C) and 6-mm Chocolate[®] balloon catheter. The CFA was ultimately dilated with a 7-mm balloon catheter (Figure 7D). A 0.014-inch CholCE PT wire was then directed into the profunda artery, and the origin was then dilated with a 5-mm balloon catheter. Finally, kissing-balloon inflations were performed in both the profunda and SFAs. The final arteriogram demonstrated wide patency of the CFA and SFA and moderate eccentric disease of the proximal profunda artery (Figure 8A). Repeat IVUS of the SFA and CFA demonstrated a significant reduction in plaque burden, calcium, and an increase in cross-sectional area (Figure 8B and 8C).

DISCUSSION

Both cases demonstrate the effectiveness of the JETSTREAM Atherectomy System in treating complex, calcified PAD. Preatherectomy IVUS characterized the location and extent of the lesion-specific calcium and the complex morphology of the disease. These cases




Figure 8. Final angiogram of the right CFA, SFA, and profunda artery (A). Final IVUS image of the right CFA (B). Final IVUS image of the right SFA origin (C).

also demonstrate that lesion morphology and complexity may not be fully appreciated by angiography alone. IVUS characterizes vessel size, extent and severity of disease, and morphologic features, including the presence and location of calcium. The JETSTREAM Atherectomy System is effective in removing plaque and calcium, resulting in significant luminal cross-sectional area. It was particularly effective for highly calcified disease of the CFA and SFA. Despite the severity and complexity of the disease in this challenging location, a favorable procedural result was achieved with combined atherectomy and balloon angioplasty.

Why Is Calcium Removal Important?

Calcium is common in patients with PAD, and its presence adversely affects procedural results and long-term outcomes. The presence of calcium necessitates greater balloon inflation pressures, resulting in an increased rate of dissections after balloon angioplasty. Despite using high balloon inflation pressures, the presence of calcium and excessive plaque burden may limit stent expansion.⁴⁻⁷ The presence of severe calcium limits the long-term effectiveness of DCBs by interfering with effective drug absorption. In a recent analysis, 12-month patency of femoropopliteal arterial segments following treatment with a DCB was 50% for lesions with calcium encompassing 270° to 360°, versus 100% for lesions with calcium from 0° to 90°. Compared to lesions with less severe calcium, excessive calcium was associated with lower ankle-brachial indices, greater late-lumen loss, and high target lesion revascularization.⁸

Benefits of the JETSTREAM Atherectomy System

The JETSTREAM Atherectomy System is effective in removing calcium in femoropopliteal disease. The safety and effectiveness of this device in removing moderate to severe superficial calcium in de novo femoral and popliteal arterial occlusive disease was evaluated in the JETSTREAM Calcium Study.⁹ This prospective, multicenter registry used IVUS before and after atherectomy to characterize the efficacy of the JETSTREAM Atherectomy System in removing calcium. The study demonstrated a significant reduction in stenosis diameter (86%, preatherectomy; 37%, postatherectomy; and 10%, postadjunct therapy) and an increase in luminal area (pre: $6.6 \pm 3.7 \text{ mm}^2$; post: $10.0 \pm 3.6 \text{ mm}^2$; $P = .001$). In addition, calcium removal was responsible for $86\% \pm 23\%$ of the increase in luminal area following treatment. The ability of the JETSTREAM Atherectomy System to remove plaque (including superficial calcium) improves luminal diameter and cross-sectional area. These luminal gains are further enhanced by adjunct balloon angioplasty (plain-old balloon angioplasty or caged balloons). By virtue of plaque modification and calcium removal, the JETSTREAM Atherectomy System may lead to improved stent results using conventional self-expanding nitinol stents.

Finally, the ability to remove calcium may also improve late results following use of adjunctive DCBs. The DEFINITIVE AR study,¹⁰ a prospective, multicenter, randomized pilot study, evaluated the use of SilverHawk™ and TurboHawk™ (Medtronic) directional atherectomy systems and Bayer HealthCare's peripheral paclitaxel-coated angioplasty catheter with Paccocath® technology. It was designed to assess the clinical benefits of plaque removal using this device, followed by treatment with a DCB with an endpoint of 12-month angiographic patency. DEFINITIVE AR demonstrated higher technical success and a lower incidence of flow-limiting dissections following this treatment strategy compared to using a DCB alone. Additionally, directional atherectomy combined with a DCB improved patency in long and severely calcified lesions. Primary patency rates for the long (> 10 cm) lesion subset at 12 months as evaluated by duplex ultrasound were 96.8% in patients treated with directional atherectomy and antirestenosis therapy (DAART) compared to 85.9% in patients treated with a DCB alone. Primary patency rates at 12 months in severely calcified lesions, per core lab assessment, were 70.4% in DAART patients,

compared to 62.5% in patients treated with DCB alone. DAART resulted in 94.1% primary angiographic patency when more plaque was removed with directional atherectomy (< 30% residual stenosis was achieved), compared to 68.8% patency when less plaque was removed (> 30% residual stenosis) before treatment with the DCB.

CONCLUSION

PAD is complex and diffuse, and it is often associated with calcium deposition. The presence of calcium may limit the effectiveness of balloon angioplasty, stenting, and DCBs. Calcium has also been associated with increased rates of dissection following balloon angioplasty. The JETSTREAM Atherectomy System is effective in removing calcium and leads to improved luminal dimensions and cross-sectional area. Plaque modification using the JETSTREAM Atherectomy System may lower complication rates (dissection) and improve early and late results using adjunctive balloon angioplasty, stenting, and/or DCBs. ■

Thomas M. Shimshak, MD, FACC, FSCAI, is an interventional cardiologist for the Heart and Vascular Center, Florida Hospital Heartland Medical Center, in Sebring, Florida. He has disclosed that he is a consultant and speaker for Bayer, Volcano Corporation, and Boston Scientific Corporation; performs training programs for Bayer and Volcano Corporation; and receives research funding from Bayer. He receives honoraria from Bayer, Volcano Corporation, and Boston Scientific Corporation. Dr. Shimshak may be reached at tshimshak@gmail.com.

1. Fowkes FG, Murray GD, Butcher I, et al. Ankle brachial index combined with Framingham risk score to predict cardiovascular events and mortality: a meta-analysis. *JAMA*. 2008;300:197-208.
2. Egorova NN, Guillermo S, Gelijns A, et al. An analysis of the outcomes of a decade of experience with lower extremity revascularization including limb salvage, lengths of stay, and safety. *J Vasc Surg*. 2010;51:878-885.
3. Shrikhande GV, Khan SZ, Hussain HG, et al. Lesion types and device characteristics that protect distal embolization during percutaneous lower extremity interventions. *J Vasc Surg*. 2011;53:347-352.
4. Bishop PD, Feiten LE, Ouriel K, et al. Arterial calcification increases in distal arteries in patients with peripheral arterial disease. *Ann Vasc Surg*. 2008;22:799-805.
5. Demer LL. Effect of calcium on in vivo mechanical response of rabbit arteries to balloon dilation. *Circulation*. 1991;83:2083-2093.
6. Makam P. Use of orbital atherectomy treatment in a high-volume clinical practice modifies non-compliant plaque to deliver durable long-term results. *J Invasive Cardiol*. 2013;25:85-88.
7. Fitzgerald PJ, Ports TA, Yock PG. Contribution of localized calcium deposits to dissection after angioplasty: an observational study using intravascular ultrasound. *Circulation*. 1992;86:64-70.
8. Fanelli F, Canavale A, Gazzetti M, et al. Calcium burden assessment and impact on drug-eluting balloons in peripheral arterial disease. *Cardiovasc Intervent Radiol*. 2014;37:898-907.
9. Akiko M, Mintz GS, Shimshak TM, et al. Intravascular ultrasound evaluation of JETSTREAM atherectomy for removal of superficial calcium in peripheral arteries. *EuroIntervention*. 2015;11:96-103.
10. Zeller T. Definitive AR. Presented at: Vascular Interventional Advances (VIVA); November 4-7, 2014; Las Vegas, Nevada.

Key Learnings From the JETSTREAM Atherectomy Calcium Study

Insights from the authors on removing severe superficial calcium to achieve significant luminal gain in femoropopliteal arteries.

BY AKIKO MAEHARA, MD; GARY S. MINTZ, MD; AND WILLIAM A. GRAY, MD

The recently published JETSTREAM Calcium Study was a prospective, single-arm, multicenter study to evaluate the effect of the JETSTREAM™ Atherectomy System (Boston Scientific Corporation) when treating severely calcified peripheral arterial lesions in the common femoral, superficial femoral, or popliteal arteries causing claudication.¹ The main question was whether the JETSTREAM Atherectomy System was effective in removing calcification. This was evaluated using both quantitative and qualitative intravascular ultrasound (IVUS), by comparing preintervention and postatherectomy IVUS images. The two major findings were as follows: The JETSTREAM Atherectomy System removed and modified moderate to severe superficial calcium to achieve significant lumen gain as standalone therapy; and adjunctive balloon angioplasty after calcium modification with the JETSTREAM Atherectomy System showed further lumen increase without major complications. In this study, the JETSTREAM 2.1/3.0 mm device was used for all procedures without distal protection. There were no major adverse events up to 30 days postprocedure.

WHY AN IVUS STUDY IS UNIQUE

Calcium was screened by angiography to identify moderate to severe obstructive intraluminal calcification in the common femoral, superficial femoral, or popliteal arteries. Lesions were evaluated by IVUS. Patients identified by angiography as possible candidates were included in the

final analysis only if there was superficial calcium that had an arc > 90° and a length > 5 mm. Overall, 55 patients were screened; however, only 26 patients met the inclusion criteria. Half of the lesions identified angiographically as having moderate to severe calcification did not have severe superfi-

CASE STUDY 1: COMMON FEMORAL*




Figure 1. Before (A) and after (B) successful revascularization of a highly stenotic left common femoral artery.




Figure 2. Pre-atherectomy IVUS image of the common femoral artery (lumen area = 2.4 mm²) (A) compared to post-JETSTREAM image (B) illustrates impressive luminal gain and a circumferential lumen created with standalone JETSTREAM Atherectomy (lumen area = 8.6 mm²). Boston Scientific images on file from the JETSTREAM Calcium Study.

...major findings were as follows:

The JETSTREAM Atherectomy System removed and modified moderate to severe superficial calcium to achieve significant lumen gain as standalone therapy.

*Results from case studies are not necessarily predictive of results in other cases. Results in other cases may vary.




Figure 3. Method of intravascular ultrasound analysis of calcium reduction. The pretreatment IVUS (A). The corresponding postatherectomy IVUS image (B). The analysis sequence is shown at the bottom. After identifying and matching the slices with calcium reduction, the lumen borders for both pretreatment (yellow circle: lumen area = 4.7 mm²) (C) and postatherectomy images (red circle: lumen area = 7.7 mm²) (E) were contoured, and the two were overlaid (D). By comparing the two contours to the visual assessment of plaque, lumen gain (3 mm²) could be attributed to a reduction of calcified plaque (blue area = 1.1 mm²) or to a reduction of noncalcified plaque (red area = 1.9 mm²). Reprinted from Maehara A, Mintz GS, Shimshak TM, et al. Intravascular ultrasound evaluation of JETSTREAM atherectomy removal of superficial calcium in peripheral arteries. *EuroIntervention*. 11(1), 96-103, Copyright 2015, with permission from Europa Digital & Publishing.

cial calcium (calcium within the lumen) at the lesion site as determined by IVUS. In these lesions, superficial calcification existed only in nonstenotic segments, or only deep calcification (calcium within the vessel wall) was present at the stenosis site. Therefore, the first finding of this study was the limitation of peripheral angiography to detect and localize calcification in peripheral arterial lesions. Deep calcification may not affect luminal gain (ie, create a stenosis). Therefore, the differentiation between superficial and deep calcification and their respective roles in severe stenosis is important when evaluating the true efficacy of any atherectomy

At the slice with the maximum calcium reduction, the lumen area increased from 6.6 ± 3.7 mm² preintervention to 10 ± 3.6 mm² ($P = .001$) after atherectomy.




Figure 4. Representative case of pretreatment, postatherectomy, and postballoon final images. The lumen increased from pretreatment (5.2 mm²) to postatherectomy (6.9 mm²) to postballoon (12.1 mm²) without dissection. Reprinted from Maehara A, Mintz GS, Shimshak TM, et al. Intravascular ultrasound evaluation of JETSTREAM atherectomy removal of superficial calcium in peripheral arteries. *EuroIntervention*. 11(1), 96-103, Copyright 2015, with permission from Europa Digital & Publishing.

procedure and device. These findings are similar to the data reported by Mintz et al in coronary artery lesions.² In that study, IVUS detected calcium in 841 of 1,155 coronary artery lesions (73%), while angiography detected calcium in only 440 (38%). Therefore, the overall sensitivity of angiography relative to IVUS was 48%, with a specificity of 89%.

SIGNIFICANT LUMINAL GAIN ACHIEVED WITH JETSTREAM ATHERECTOMY

For the patients who were ultimately included in the study, first the preintervention and postatherectomy IVUS lumens were outlined. Second, the postatherectomy IVUS images were overlaid onto their respective preintervention images. Assuming there was no change in total arterial area, the change in lumen area was attributed to either calcified plaque or noncalcified plaque removal (Figure 3). At the slice with the maximum calcium reduction, the lumen area increased from 6.6 ± 3.7 mm² preintervention to 10 ± 3.6 mm² ($P = .001$) after atherectomy. The decrease in calcium area, measured as 2.8 ± 1.6 mm², was responsible for $86\% \pm 23\%$ of the lumen area increase. Additionally, the arc of reverberations increased from 25° (range, 15°–35°) to 70° (range, 46°–95°), $P = .001$, indicating device-related modification of calcium. Therefore, the second lesson was that the JETSTREAM Atherectomy System increased lumen dimensions by calcium removal as well as by calcium modification (increase in reverberations).

CASE STUDY 2: DISTAL SFA/PROXIMAL POPLITEAL*




Figure 5. Successful debulking with the JETSTREAM Atherectomy System in a distal right SFA/proximal popliteal artery lesion (A). The pre-atherectomy IVUS image (B) reveals a lumen area of 2.5 mm². The post-atherectomy images (C and D) reveal a lumen area of 7.6 mm² and impressive debulking with JETSTREAM Atherectomy even before adjunctive therapy. Boston Scientific images on file from the JETSTREAM Calcium Study.

VESSEL EXPANSION WITHOUT VESSEL DAMAGE

In the 11 lesions that had postadjunctive balloon IVUS images, the minimum lumen area increased further from 7 mm² (range, 6.4–7.8 mm²) after atherectomy to 11.9 mm² (range, 10.3–13.5 mm²) after adjunct balloon inflation ($P < .01$). However, the prevalence of dissections also increased from 3/11 after atherectomy to 8/11 after adjunct balloon inflations ($P = .03$). However, the maximum angle of the dissection flap was minor (42° [range, 17°–66°]) with a preserved lumen area (15.6 mm² [range, 13.4–17.7 mm²]) within the dissection. The dissections were non-flow limiting. Also,

the higher resolution of IVUS imaging versus angiography most likely led to a higher detection rate. Thus, the third and final lesson was that the JETSTREAM Atherectomy System allowed additional lumen increase by facilitating vessel expansion without significant vessel damage (ie, dissection), presumably because of calcium modification. A representative case is shown in Figure 4.

CONCLUSION

Severely calcified lesions may cause damage to the polymer/drug coating of a drug-eluting stent, resulting in inadequate drug delivery.^{3,4} Although there is accumulating evidence in coronary artery intervention showing that calcified lesions have worse outcomes compared to noncalcified lesions,^{5,6} the clinical impact of superficial calcium removal in peripheral artery disease in respect to effectiveness of drug-coated balloons or drug-eluting stents needs further investigation.⁷ ■

Akiko Maehara, MD, is with Columbia University Medical Center and the Cardiovascular Research Foundation in New York, New York. Dr. Maehara has disclosed that she receives a grant from and is a consultant to, Boston Scientific Corporation.

Gary S. Mintz, MD, is with the Cardiovascular Research Foundation in New York, New York. Dr. Mintz has disclosed that he receives grant support and is a consultant for Boston Scientific Corporation and Volcano Corporation.

William A. Gray, MD, is with Columbia University Medical Center in New York, New York. He has disclosed that he is a consultant to Boston Scientific Corporation. Dr. Gray may be reached at (212) 304-5697; wg2131@columbia.edu.

Acknowledgement: The authors thank investigators Thomas M. Shimshak, MD; Joseph J. Ricotta II, MD, MS; Venkatesh Ramaiah, MD; Malcolm T. Foster III, MD; and Thomas P. Davis, MD, for their work on the JETSTREAM Calcium Study.

1. Maehara A, Mintz GS, Shimshak TM, et al. Intravascular ultrasound evaluation of JETSTREAM atherectomy removal of superficial calcium in peripheral arteries. *EuroIntervention*. 2015;11:96–103.
2. Mintz GS, Popma JJ, Pichard AD, et al. Patterns of calcification in coronary artery disease. A statistical analysis of intravascular ultrasound and coronary angiography in 1155 lesions. *Circulation*. 1995;91:1959–1965.
3. Hwang CW, Edelman ER. Arterial ultrastructure influences transport of locally delivered drugs. *Circ Res*. 2002;90:826–832.
4. Kuriyama N, Kobayashi Y, Yamaguchi M, et al. Usefulness of rotational atherectomy in preventing polymer damage of everolimus-eluting stent in calcified coronary artery. *JACC Cardiovasc Interv*. 2011;4:588–589.
5. Généreux P, Madhavan MV, Mintz GS, et al. Ischemic outcomes after coronary intervention of calcified vessels in acute coronary syndromes. Pooled analysis from the HORIZONS-AMI (Harmonizing Outcomes With Revascularization and Stents in Acute Myocardial Infarction) and ACUTY (Acute Catheterization and Urgent Intervention Triage Strategy) TRIALS. *J Am Coll Cardiol*. 2014;63:1845–1854.
6. Bourantas CV, Zhang YJ, Garg S, et al. Prognostic implications of coronary calcification in patients with obstructive coronary artery disease treated by percutaneous coronary intervention: a patient-level pooled analysis of 7 contemporary stent trials. *Heart*. 2014;100:1158–1164.
7. Siablis D, Kitrou PM, Spiliopoulos S, et al. Paclitaxel-coated balloon angioplasty versus drug-eluting stenting for the treatment of infrapopliteal long-segment arterial occlusive disease: the IDEAS randomized controlled trial. *JACC Cardiovasc Interv*. 2014;7:1048–1056.

*Results from case studies are not necessarily predictive of results in other cases. Results in other cases may vary.

Peripheral Rotablator™ Atherectomy: The Below-the-Knee Approach to Address Calcium Head On

Peripheral Rotablator's front-cutting, diamond-tipped burr provides stable rotation in calcified lesions.

BY SONYA S. NOOR, MD, FACS

There are multiple endovascular options for treatment of infrainguinal disease, but treatment of severe calcific disease of the superficial femoral artery (SFA), popliteal artery, and tibial vessels remains a challenge. Peripheral atherectomy is a unique treatment modality because it allows debulking of plaque with luminal gain and minimal barotrauma. This results in less injury to the vessel during initial treatment and theoretically reduces hyperplastic reaction to the initial treatment. In severely calcific vessels, calcium debulking changes the vessel wall compliance with the removal of calcium. It can then be treated with low-pressure balloon inflation with minimal injury to the vessel wall. This is now a particularly attractive concept with the availability of drug-coated balloons and drug-eluting stents, as the vessel can be prepared with atherectomy before delivery of these devices. This may ensure adequate drug delivery to the tissue, thereby reducing intimal hyperplastic reaction and increasing durability of

the procedures. Prevailing concerns with atherectomy (ie, dissection, perforation, clinically significant embolization, and durability) have prevented the widespread use of atherectomy.¹

The Peripheral Rotablator Rotational Atherectomy System (Boston Scientific Corporation) (Figure 1) is one of the newer additions to the peripheral atherectomy device field. The coronary Rotablator Atherectomy system has been used for the last 20 years, and it has been very successful in treating moderate and severe calcific coronary disease, with encouraging safety and efficacy data to support its use.² Our center has been one of the largest users of Coronary Rotablator Atherectomy, so when the Peripheral Rotablator Atherectomy System became available, we quickly adopted this technology to use in severely calcific vessels. We started using the Rotablator Atherectomy System to treat severe tibial vessel calcification and small popliteal vessels. As our experience grew, we then started to use Rotablator Atherectomy more broadly when treating




Figure 1. The Peripheral Rotablator Atherectomy System advancer (A), console (B), pedal (C), and burr (D).

The Rotablator Atherectomy System's front-cutting diamond burr is very useful in severe calcific stenotic lesions, as it can ablate its way through the calcium and create a channel that is smooth and has a predictable concentric lumen.

infrainguinal calcific disease, and found a reduction in the use of stents (or only focal stenting was required).

ROTABLATOR FEATURES AND MECHANISM OF ACTION

The Peripheral Rotablator Atherectomy System is fairly simple to use and requires a connection to the console, a power source, a compressed nitrogen tank, and an IV fluid mix to start using the device. The foot pedal starts the rotational atherectomy. Usually, 20- to 30-second runs are done under live fluoroscopy. This device uses a front-cutting diamond-coated burr to ablate the calcium particles, while rotating at 160,000 to 180,000 RPM in the lumen of the vessel. The coronary literature has studied the ablation particles over the last 20 years, and when proper technique is employed, the ablation particles generated measured about 5 μm , which is smaller than a red blood cell. These particles are then washed downstream during the treatment and picked up by the reticular endothelial system. For this reason, embolic protection devices are of no use, as the pore size of embolic protection devices are generally in the range of 100 μm and would not catch the ablation particles. Rotablator Atherectomy can only be performed with a 0.009-inch wire, which does not support the use of embolic protection devices, either.




The Rotablator Atherectomy System's front-cutting diamond burr is very useful in moderately and severe calcific stenotic lesions, as it can ablate its way through the calcium and create a channel that is smooth and has a predictable concentric lumen.

Other devices have a leading edge, which has to be introduced first through the lumen before the device can be introduced and treatment can be performed, requiring predilatation or dottering, and causing barotrauma to the vessel prior to treatment. We have found this front-cutting feature particularly helpful in moderate and severe calcific disease and when negotiating even a predilatation balloon catheter can prove difficult.

As the diamond-coated burr engages the lesion while it rotates on the RotaWire™ (Boston Scientific Corporation), it has a stable circular rotation that creates a smooth, predictably concentric lumen. It ablates the plaque with predictable ablation particles with minimal injury to the vessel wall, which can be a concern with other atherectomy devices in small vessels. Average Rotablator™ Atherectomy run times, even in

ABLATION TECHNIQUE: COMMON CONSIDERATIONS

- 160,000 to 180,000 RPM setting is optimal for above- and below-the-knee calcific lesions
- Run for 20 to 30 seconds under live fluoroscopy
- One-burr approach is common
- Limit RPM drop to under 5,000 RPM
- Plaque modification: burr-to-artery ratio, 70%–85% to native lumen diameter



(Images courtesy of Boston Scientific Corporation.)

long, diffuse lesions, are typically 3 to 4 minutes per vessel, making this an efficient treatment modality.

ROTABLATOR BEST PRACTICES

As with other atherectomy devices, the Rotablator Atherectomy System has its own learning curve and performs well when proper technique is employed. We have used the Rotablator™ Atherectomy System in severe calcific disease as the first line of treatment for these lesions. We currently use it in the SFA, popliteal, and tibial vessels as the first line of treatment. Rotablator Atherectomy proves to be an effective tool for calcific ablation requiring low atmospheric balloon postdilatation and only focal stenting, if at all necessary. We are also starting to use Rotablator Atherectomy for vessel preparation before drug-coated balloon usage in order to potentially improve drug uptake in the vessel wall.

We have found that the 160,000 to 180,000 RPM setting is optimal for both above- and below-knee calcific lesions. Twenty- to 30-second runs under live fluoroscopy and a slow, deliberate pecking action with the burr engaging the lesion for a second or two, followed by a gentle pullback, results in successful luminal gain. It is important to engage the lesion with the burr, but also pull back for 1 to 2 seconds, which allows dissipation of frictional heat and flushing of microparticles into the distal circulation. Overzealous advancement of the burr can lead to the device stalling within calcific disease and distal embolization, which should be avoided.

CASE 1*

A 71-year-old African American woman with end-stage renal disease, a previous cerebrovascular accident, coronary artery disease, and coronary artery bypass




Figure 2. A diagnostic angiogram showing severe stenotic disease of the posterior tibial artery (arrow) and complete occlusion of the anterior tibial artery (bracket) with distal reconstitution.




Figure 3. A diagnostic angiogram showing complete occlusion of the anterior tibial artery with distal reconstitution (arrow) and no flow in the distal posterior tibial artery.




Figure 4. Angiogram after posterior tibial Rotablator Atherectomy showing healthy flow in the posterior tibial artery.



Figure 5. After Rotablator Atherectomy was performed in the posterior tibial and anterior tibial arteries, healthy flow was seen in both vessels down to the foot.

*Results from case studies are not necessarily predictive of results in other cases. Results in other cases may vary.

TABLE 1. ROTABLATOR ATHERECTOMY SYSTEM SIZING

Burr (mm)	Diameter (inches)	Minimum Recommended Introducer/Guide Sheath Size (F)
1.25	0.049	4 / 5*
1.50	0.059	5
1.75	0.069	6
2.00	0.079	6 / 7*
2.25	0.089	7
2.50	0.098	7 / 8*

** For a given French size guide sheath or introducer sheath, the internal lumen and hemostasis valve will vary from manufacturer to manufacturer. When using an introducer for the first time, it should be tested with the largest Peripheral RotaLink Plus burr intended to be used with it.*

graft surgery presented with a left heel ulcer that was not healing despite treatment for many months. The patient's arterial Doppler showed noncompressible vessels and a toe-brachial index of 0.1, with reduced waveforms at all levels.

Procedural Details

Left common femoral artery ultrasound and puncture was performed with placement of a right side sheath, and a diagnostic angiogram was obtained, which clearly showed complete occlusion of the posterior tibial and anterior tibial artery (Figures 2 and 3). The patient underwent placement of a 5-F, 70-cm sheath and was heparinized before successful crossing of the posterior tibial artery occlusion and subsequently, the anterior tibial artery. The patient underwent Rotablator Atherectomy with a 1.75-mm burr of the posterior tibial artery first (Figure 4), followed by the anterior tibial artery. The total run time was 4 minutes, and the 180,000 RPM setting

was used in both arteries. After atherectomy was completed, low-pressure balloon angioplasty was performed using a 2.5-mm X 220-mm balloon, for a total of 2 minutes for each inflation.

Completion angiography showed no evidence of dissection, perforation, or distal embolization (Figure 5).

At 5-month follow-up, the patient was found to have almost completely healed ulcers, and arterial Dopplers showed improved waveforms at all levels, noncompressible ankle-brachial indices (ABIs), and a toe-brachial index of 0.5.

CASE 2*

A 65-year-old man presented with severe claudication and ischemic ulceration of the right second toe. Arterial Doppler exam showed an ABI of 0.58 on the right and 0.90 on the left. Pulmonary vascular resistance waveforms indicated distal SFA and popliteal artery disease.

Procedural Details

The patient underwent left common femoral artery access with placement of a 5-F sheath. Diagnostic angiography confirmed complete occlusion of the right SFA and popliteal artery at the adductor canal (Figure 6), with reconstitution of a popliteal artery at the knee joint and some mild diffuse tibial vessel disease. The patient underwent placement of a 7-F, 70-cm sheath, after which the patient was heparinized. A stiff Glidewire (Terumo Interventional Systems) and a 0.035-inch Quick-Cross catheter (Spectranetics Corporation) was used to cross the total occlusion, and reentry into the popliteal artery was confirmed. The RotaWire was placed into the tibial vessels, and Rotablator Atherectomy was performed using a 2.5-mm burr. A setting of 170,000 to 180,000 RPM was used for a total of 5 minutes. Then a 5- X 220-mm-long Sterling™ Balloon (Boston Scientific Corporation) was used for low-pressure angioplasty of 4 atm, for a total of 3 minutes. A follow-up angiogram (Figure 7) revealed excellent



Figure 6. A diagnostic angiogram showing occlusion of the distal SFA and popliteal artery.



Figure 7. Angiogram after Rotablator Atherectomy and low-pressure angioplasty. Restoration of flow is noted in the SFA and popliteal artery.

*Results from case studies are not necessarily predictive of results in other cases. Results in other cases may vary.

flow, good luminal gain, and no evidence of perforation, dissection, or significant embolization.

The patient was seen in follow-up 1 month after the procedure with no complaints of claudication, near complete ulceration healing, and arterial Doppler exams that showed an ABI of 0.84 on the right and 0.77 on the left, with good waveforms at all levels.

CONCLUSION

The Rotablator Atherectomy System has been used to treat moderate and severe calcific disease safely and efficiently for over 20 years in the coronary vasculature, and we started to use Rotablator Atherectomy to treat similar calcific disease in the periphery. At our center, we now use Rotablator Atherectomy as the first line of treatment whenever we encounter moderate or severe calcific disease. We have found the Rotablator Atherectomy System to be easy to set up and use, and it is efficient in ablating and treating calcium with short procedure times. There have been minimal dissections, perforation, or clinically significant embolization. As with all atherectomy devices, it is important to use proper technique while handling the device to minimize complications. The benefit of the front-cutting diamond burr is

especially useful in negotiating tight stenotic or occlusive lesions (where no predilatation is necessary), thereby minimizing barotrauma to the vessel before treatment. The stable rotation of the burr engages the calcium and ablates it, leaving a predictable concentric lumen. We have been successful in changing vessel compliance with calcium ablation, allowing minimal adjunctive therapy (such as low-pressure angioplasty, no stenting, or only focal stenting). We are also starting to use the Rotablator Atherectomy System to remove the calcium plaque burden and prep the vessel before use of drug-coated balloons. It remains to be seen if this strategy enhances drug uptake into the vessel wall, and therefore increase the durability of this procedure. ■

Sonya S. Noor, MD, FACS, is a clinical associate professor with the University at Buffalo Gates Vascular Institute in Buffalo, New York. She has disclosed that she is a consultant for Boston Scientific Corporation. Dr. Noor may be reached at snoor@kaleidahealth.org.

1. Zeller T, Frank U, Bürgelin K, et al. Initial clinical experience with percutaneous atherectomy in the infragenicular arteries. *J Endovasc Ther.* 2003;10:987-993.

2. Reisman MD. *Guide to Rotational Atherectomy*. Birmingham, MI: Physicians Press; 1997.

(Continued from page 7)

in very heavily calcified occlusions and long lesions, you can get good results with patency rates of 90% at 1 year.

EVT: In your experience, what is the best way to remove calcium from these particular lesions?

Prof. van den Berg: I think you need some kind of mechanical atherectomy. I'm using laser atherectomy a lot, but in these cases merely as a tool to modify the calcium because we know that laser is not really good at completely removing calcium.

EVT: Are there particular characteristics of the balloons that you use for percutaneous transluminal angioplasty that you find to be advantageous?

Prof. van den Berg: I typically use semicompliant balloons that give me a little bit of space to play around with the diameter. By using the compliance chart, the diameter of the balloon can be adapted to the diameter of the vessel wall. It's very important to be aware of the fact that when you use compliant or semicompliant balloons when there is a tight stenosis, in the areas where the balloon opens up more than in the area of the tight stenosis, the vessel wall might get injured much more at the proximal and distal end of the balloon (ie, the dog-bone effect). That might, again, be a factor that is influencing restenosis in the long term. ■

1. Fanelli F, Cannavale A, Gazzetti M, et al. Calcium burden assessment and impact on drug-eluting balloons in peripheral arterial disease. *Cardiovasc Intervent Radiol.* 2014;37:898-907.

2. Cioppa A, Stabile E, Popuso G, et al. Combined treatment of heavy calcified femoro-popliteal lesions using directional atherectomy and a paclitaxel coated balloon: one-year single centre clinical results. *Cardiovasc Revasc Med.* 2012;13:219-223.

Results from case studies are not necessarily predictive of results in other cases. Results in other cases may vary.

ABBREVIATED STATEMENTS

JETSTREAM CATHETERS COMBINED WITH CONSOLE

CAUTION: Federal law (USA) restricts this device to sale by or on the order of a physician. Rx only. Prior to use, please see the complete "Instructions for Use" for more information on Indications, Contraindications, Warnings, Precautions, Adverse Events, and Operator's Instructions.

Catheter INDICATIONS

The Jetstream System is intended for use in atherectomy of the peripheral vasculature and to break apart and remove thrombus from upper and lower extremity peripheral arteries. It is not intended for use in coronary, carotid, iliac or renal vasculature.

Console INDICATIONS

The PV Console is designed for use only with the Jetstream Catheter and Control Pod. See the current revision of the applicable Catheter and Control Pod Instructions for Use for further information.

CONTRAINDICATIONS

No known contraindications.

Catheter WARNINGS/PRECAUTIONS

- The Jetstream Catheter and Control Pod may only be used with the PV Console.
- Take care to avoid being pinched when closing the aspiration and infusion pump doors. Use room temperature infusate only. Use of heated infusate may lead to wrinkling, ballooning and/or bursting of the outer catheter sheath.
- Do not bend or kink the Catheter during setup or during the procedure. This may damage the device and lead to device failure.
- Operating the Catheter over a kinked guidewire may cause vessel damage or guidewire fracture.
- During treatment, do not allow the Catheter tip within 10.0 cm of spring tip portion of the guidewire. Interaction between the Catheter Tip and this portion of the guidewire may cause damage to or detachment of the guidewire tip or complicate guidewire management.
- The guidewire must be in place prior to operating the Catheter in the patient. Absence of the guidewire may lead to inability to steer the Catheter and cause potential vessel damage.
- Do not inject contrast while the device is activated.
- If the guidewire is accidentally retracted into the device during placement or treatment, stop use, and remove the Catheter and the guidewire from the patient. Verify that the guidewire is not damaged before re-inserting the guidewire. If damage is noticed, replace the guidewire.
- Check the infusate bag frequently and replace when needed. Do not run the JETSTREAM System without infusate as this may cause device failure.
- Hold the guidewire firmly during Catheter retraction process. Failure to do so may result in guidewire rotation within the vessel.
- Do not manipulate the Catheter against resistance unless the cause for that resistance has been determined.
- Use only listed compatible guidewires and introducers with the Jetstream System. The use of any supplies not listed as compatible may damage or compromise the performance of the Jetstream System.

Prior to use of the Jetstream System, confirm the minimum vessel diameter proximal to the lesion per the following:

Jetstream SC Atherectomy Catheter 1.6 Minimum Vessel Diameter Proximal to Lesion 2.5 mm

Jetstream SC Atherectomy Catheter 1.85 Minimum Vessel Diameter Proximal to Lesion 2.75 mm

Jetstream XC Atherectomy Catheter 2.1-3.0 Minimum Vessel Diameter, Blades Down 3.0 mm; Minimum Vessel Diameter, Blades Up 4.0 mm

Jetstream XC Atherectomy Catheter 2.4-3.4 Minimum Vessel Diameter, Blades Down 3.5 mm; Minimum Vessel Diameter, Blades Up 4.5 mm

Console WARNINGS/PRECAUTIONS

- WARNING: To avoid the risk of electric shock, this equipment must only be connected to a supply mains with protective earth.
- Do not open either pump door during operation of the System. Doing so could result in loss of aspiration and/or infusion and will halt device activation.
- Ensure the PV Console display is visible during the entire procedure.
- Observe normal safety practices associated with electrical/electronic medical equipment.
- Avoid excessive coiling or bending of the power cables during storage.
- Store the PV Console using appropriate care to prevent accidental damage.
- Do not place objects on the PV Console.
- Do not immerse the PV Console in liquids.

ADVERSE EVENTS

Potential adverse events associated with use of this device and other interventional catheters include, but are not limited to the following (alphabetical order):

- Abrupt or sub-acute closure • Amputation • Bleeding complications, access site • Bleeding complications, non-access site • Death • Dissection • Distal emboli • Hypotension • Infection or fever • Perforation • Restenosis of the treated segment • Vascular complications which may require surgical repair • Thrombus • Vasospasm

ROTABLATOR PERIPHERAL ROTALINK PLUS

ROTABLATOR PERIPHERAL ROTAWARE GUIDEWIRE AND WIRECLIP TORQUER

ROTABLATOR ROTATIONAL ATHERECTOMY SYSTEM CONSOLE

ROTAGLIDE LUBRICANT

CAUTION: Federal law (USA) restricts this device to sale by or on the order of a physician. Rx only. Prior to use, please see the complete "Directions for Use" for more information on Indications, Contraindications, Warnings, Precautions, Adverse Events, and Operator's Instructions.

Rotalink Plus INTENDED USE/INDICATIONS FOR USE

The Rotablator Rotational Atherectomy System is intended for percutaneous use in the peripheral vessels in patients with occlusive atherosclerotic disease who are acceptable candidates for endovascular procedures.

RotaWire: INDICATIONS FOR USE/INTENDED USE

These guidewires are intended for use with the Rotablator Rotational Atherectomy System.

Lubricant INDICATIONS FOR USE

Rotaglide lubricant is intended for use with the Rotablator atherectomy system, for the purpose of increasing the lubricity of the system.

CONTRAINDICATIONS AND RESTRICTIONS

Contraindications

1. Occlusions through which a guidewire will not pass.
2. Use in coronary arteries.
3. Long (≥ 20 cm) total occlusions.
4. Angiographic evidence of thrombus prior to treatment with the Rotablator Rotational Atherectomy System. Such patients may be treated with thrombolytics (e.g., Urokinase). When the thrombus has been resolved for two to four weeks, the lesion may be treated with the Rotablator Rotational Atherectomy System.
5. Angiographic evidence of significant dissection at the treatment site. The patient may be treated conservatively for approximately four weeks to permit the dissection to heal before treating the lesion with the Rotablator Rotational Atherectomy System.

Lubricant CONTRAINDICATIONS

Rotaglide™ lubricant is contraindicated in patients with known allergies to the lubricant ingredients: olive oil, egg yolk phospholipids, glycerin, sodium deoxycholate, L-histidine, disodium EDTA, sodium hydroxide, and water.

Restrictions

- Federal (USA) law restricts the use of this system to physicians who are credentialed in peripheral angioplasty and who have attended the Rotablator System Physician Training Program.

WARNINGS

- The risks of Rotational Atherectomy can be reduced if the device and associated accessories are used in the appropriate patient population by a physician who has had adequate training.
- If the Peripheral RotaLink Plus shows evidence of mechanical failure at any time prior to or during the angioplasty procedure, immediately discontinue use of the device and return it to Customer Service for evaluation. Do NOT attempt to use a damaged Peripheral RotaLink Plus; use may result in device malfunction and/or patient injury.
- Never operate the Peripheral RotaLink Plus without saline infusion. Flowing saline is essential for cooling and lubricating the working parts of the advancer. Operation of the advancer without proper saline infusion may result in permanent damage to the advancer.
- Never operate the Peripheral RotaLink Plus with the Rotablator Rotational Atherectomy System in Dynaglide™ mode or operate the guidewire brake defeat button unless you have a firm grip on the guidewire using the wireClip™ Torquer. The wireClip Torquer may be held with the fingers or inserted com-

pletely into the docking port after the brake button is depressed. Defeating the brake, or operating the Peripheral RotaLink Plus with the Rotablator Rotational Atherectomy System in Dynaglide mode, without securing the guidewire may result in rotation and entanglement of the guidewire.

- During setup of the Peripheral RotaLink Plus never grip or pull on the flexible shaft.
- The burr at the distal tip of the Peripheral RotaLink Plus is capable of rotating at very high speeds. Do NOT allow parts of the body or clothing to come in contact with the burr. Contact may result in physical injury or entanglement.
- Never advance the rotating burr to the point of contact with the guidewire spring tip. Such contact could result in distal detachment and embolization of the tip.
- If the Peripheral RotaLink Plus stops and the red STALL light on the console illuminates, retract the burr and immediately discontinue treatment. Check the advancer for proper connection to the console. If the connections are correct, use fluoroscopy to analyze the situation. Never force the system when rotational or translational resistance occurs, as vessel perforation may occur.
- Never advance the rotating burr by advancing the sheath. Guidewire buckling may occur and perforation or vascular trauma may result. Always advance the rotating burr by using the advancer knob.
- If resistance is encountered, retract the burr and stop treatment immediately. Use fluoroscopy to analyze the situation. Never force the Peripheral RotaLink Plus when rotational or translational resistance occurs, as vessel perforation, vessel trauma or embolism due to burr detachment or fractured wire may occur and in rare instances may result in surgical intervention and death.
- The use of Rotablator Rotational Atherectomy System for in-stent stenosis might lead to damage of stent components and/or Peripheral RotaLink Plus, which may lead to patient injury.
- Always keep the burr advancing or retracting while it is rotating. Maintaining the burr in one location while it is rotating may lead to excessive tissue removal or damage to the Peripheral RotaLink Plus or entrapment of the Peripheral RotaLink Plus. It is best to advance and retreat the burr no more than 3 cm at a time in a smooth pecking motion, being careful to engage the lesion only minimally when resistance is met. Do not allow the individual burr run time to exceed 30 seconds with total rotational procedure time not to exceed five minutes.

RotaWire WARNINGS

Use extreme caution and careful judgment in patients for whom anticoagulation is not indicated. Console WARNINGS

- Never use oxygen as the propellant for the Rotablator Rotational Atherectomy System.
- The use of accessories, transducers and cables other than those specified, with the exception of transducers and cables sold by the manufacturer of the Rotablator System as replacement parts for internal components, may result in increased emissions or decreased immunity of the Rotablator System.
- This device is not to be used in the presence of flammable anesthetics.
- Do NOT operate the Rotablator Console with gas pressures in excess of 758.4 kPa (110 psi).
- Do not modify or repair.

Lubricant WARNINGS

Discard vial if there are particulates in the emulsion or if an oiling-out of emulsion has occurred.

PRECAUTIONS

- Percutaneous rotational angioplasty with the Rotablator Rotational Atherectomy System should only be carried out at hospitals where emergency bypass surgery can be immediately performed in the event of a potentially injurious or life-threatening complication.
- Appropriate drug therapy including (but not limited to) anticoagulant and vasodilator therapy must be provided to the patient during all phases of patient care.
- When the Peripheral RotaWire™ Guidewires and/or Peripheral RotaLink Plus are in the body, they should only be manipulated while they are under fluoroscopic observation with radiographic equipment that provides high resolution images.
- Use only normal saline as the infusate. Never inject contrast agent, or any other substance that is not approved as part of the Rotablator Rotational Atherectomy System, into the infusion port or saline infusion bag as this may cause permanent damage to the Peripheral RotaLink Plus.

Console PRECAUTIONS

- User should take precautions when using the console in conjunction with other medical electrical equipment.
- The Rotablator Console needs special precautions regarding EMC and needs to be installed and put into service according to the EMC information provided in Appendix D in the DFU.

ADVERSE EVENTS

Potential adverse reactions which may result from the use of this device include but are not limited to:

- Additional intervention • Allergic reaction • Amputation • Death • Embolism • Hematoma/Hemorrhage • Hemodynamic changes • Hemoglobinuria • Infection • Restenosis • Stroke • Slow, no flow, abrupt vessel closure • Surgery including arterial bypass • Thrombosis and vessel occlusion • Vessel trauma (dissection, perforation, pseudoaneurysm, arteriovenous fistula)
- There may also be complications associated with distortion, kinks, and fracture of the guidewire and physical deterioration or malfunction of the device, which can lead to patient injury or death.

SOLENT CATHETERS COMBINED W/CONSOLE

SOLENT OMNI, SOLENT PROXI THROMBECTOMY CATHETERS

SOLENT DISTA THROMBECTOMY CATHETER

ANGIOJET ULTRA CONSOLE

CAUTION: Federal law (USA) restricts this device to sale by or on the order of a physician. Rx only. Prior to use, please see the complete "Instructions for Use" for more information on Indications, Contraindications, Warnings, Precautions, Adverse Events, and Operator's Instructions.

INDICATIONS AND USAGE

The Angiojet SOLENT proxi & omni Thrombectomy Sets are intended for use with the Angiojet Ultra Console to break apart and remove thrombus from:

- upper and lower extremity peripheral arteries ≥ 3.0 mm in diameter,
 - upper extremity peripheral veins ≥ 3.0 mm in diameter,
 - iliofemoral and lower extremity veins ≥ 3.0 mm in diameter,
 - A-V access conduits ≥ 3.0 mm in diameter and
 - for use with the Angiojet Ultra Power Pulse technique for the control and selective infusion of physician specified fluids, including thrombolytic agents, into the peripheral vascular system.
- The Angiojet SOLENT dista Thrombectomy Set is intended for use with the Angiojet Ultra Console to break apart and remove thrombus from:
- upper and lower extremity peripheral arteries and
 - for use with the Angiojet Ultra Power Pulse technique for the control and selective infusion of physician specified fluids, including thrombolytic agents, into the peripheral vascular system.

The minimum vessel diameter for each Thrombectomy Set model is listed in Table 1 (in the IFU).

CONTRAINDICATIONS

Do not use the catheter in patients:

- Who are contraindicated for endovascular procedures
- Who cannot tolerate contrast media
- In whom the lesion cannot be accessed with the guide wire

WARNINGS AND PRECAUTIONS

- The Thrombectomy Set has not been evaluated for treatment of pulmonary embolism. There are reports of serious adverse events, including death, associated with cases where the catheter was used in treatment of pulmonary embolism.
- The Thrombectomy Set has not been evaluated for use in the carotid or cerebral vasculature.
- The Thrombectomy Set has not been evaluated for use in the coronary vasculature (unless accompanied by a separate coronary IFU).
- Operation of the catheter may cause embolization of some thrombus and/or thrombotic particulate debris. Debris embolization may cause distal vessel occlusion, which may further result in hypoperfusion or tissue necrosis.
- Cardiac arrhythmias during catheter operation have been reported in a small number of patients. Cardiac rhythm should be monitored during catheter use and appropriate management, such as temporary pacing, be employed, if needed.
- Use of the catheter may cause a vessel dissection or perforation.
- Do not use the Angiojet Ultra System in patients who have a nonhealed injury due to recent mechanical intervention, in the vessel to be treated, to avoid further injury, dissection, or hemorrhage.
- Do not use the Thrombectomy Set in vessels smaller than minimum vessel diameter for each Thrombectomy Set model as listed in Table 1 (in the IFU); such use may increase risk of vessel injury.
- Systemic heparinization is advisable to avoid pericatheterization thrombus and acute rethrombosis. This is in addition to the heparin added to the saline supply bag.

- Operation of the AngioJet System causes transient hemolysis which may manifest as hemoglobinuria. Table 1 (in the IFU) lists maximum recommended run times in a flowing blood field and total operating time for each Thrombectomy Set. Evaluate the patient's risk tolerance for hemoglobinemia and related sequelae prior to the procedure. Consider hydration prior to, during, and after the procedure as appropriate to the patient's overall medical condition.
- Large thrombus burdens in peripheral veins and other vessels may result in significant hemoglobinemia which should be monitored to manage possible renal, pancreatic, or other adverse events.
- Monitor thrombotic debris/fluid flow exiting the Thrombectomy Set via the waste tubing during use. If blood is not visible in the waste tubing during AngioJet Ultra System activation, the catheter may be occlusive within the vessel; verify catheter position, vessel diameter and thrombus status. Operation under occlusive conditions may increase risk of vessel injury.
- Do not exchange the guide wire. Do not retract the guide wire into the catheter during operation. The guide wire should extend at least 3 cm past the catheter tip at all times. If retraction of the guide wire into the Thrombectomy Set occurs, it may be necessary to remove both the Thrombectomy Set and the guide wire from the patient in order to re-load the catheter over the guide wire. (Distal only)
- Use of a J-tip guide wire is not recommended as it is possible for the tip of the guide wire to exit through a side window on the distal end of the catheter. (Omni, Proxi only)
- Do not pull the catheter against abnormal resistance. If increased resistance is felt when removing the catheter, remove the catheter together with the sheath or guide catheter as a unit to prevent possible tip separation.
- If resistance is felt during the advancement of the Thrombectomy Set to lesion site, do not force or torque the catheter excessively as this may result in deformation of tip components and thereby degrade catheter performance.
- Obstructing lesions that are difficult to cross with the catheter to access thrombus may be balloon dilated with low pressure (≤ 2 atm). Failure to pre-dilate difficult-to-cross lesions prior to catheter operation may result in vessel injury.
- The potential for pulmonary thromboembolism should be carefully considered when the Thrombectomy Sets are used to break up and remove peripheral venous thrombus.
(Below is Omni, Proxi only)
- Hand injection of standard contrast medium may be delivered through the thrombectomy catheter via the manifold port stopcock. Follow the steps to remove air from the catheter when delivering fluid through the catheter stopcock.
- Fluids should be injected only under the direction of a physician and all solutions prepared according to the manufacturer instructions.
- The Thrombectomy Set waste lumen is rated for 50psi. Delivering a hand injection of contrast medium with excessive force can create injection pressures greater than 50psi, potentially causing leaks in the waste lumen of the catheter.
- Do not use a power injector to deliver contrast medium through the catheter stopcock. Power injectors can deliver pressures greater than 50psi, potentially causing leaks in the waste lumen of the catheter.
- Some fluids, such as contrast agents, can thicken in the catheter lumen and block proper catheter operation if left static too long. The catheter should be operated to clear the fluid within 15 minutes of injection.

Console WARNINGS and PRECAUTIONS:

- Use the AngioJet Ultra Console only with an AngioJet Ultra Thrombectomy Set. This Console will not operate with a previous model pump set and catheter.
- Do not attempt to bypass any of the Console safety features.
- If the catheter is removed from the patient and/or is inoperative, the waste tubing lumen, guide catheter, and sheath should be flushed with sterile, heparinized solution to avoid thrombus formation and maintain lumen patency. Reprime the catheter by submerging the tip in sterile, heparinized solution and operating it for at least 20 seconds before reintroduction to the patient.
- Refer to the individual AngioJet Ultra Thrombectomy Set Information for Use manual for specific warnings and precautions.
- Do not move the collection bag during catheter operation as this may cause a collection bag error.
- Monitor thrombotic debris/fluid flow exiting the catheter through the waste tubing during use. If blood is not visible during console activation, the catheter may be occlusive within the vessel or the outflow lumen may be blocked.
- Ensure adequate patient anticoagulation to prevent thrombus formation in outflow lumen.
- Refer to individual Thrombectomy Set Instructions for Use manual for specific instructions regarding heparinization of the Thrombectomy Set.
- The Console contains no user-serviceable parts. Refer service to qualified personnel.
- Removal of outer covers may result in electrical shock.
- This device may cause electromagnetic interference with other devices when in use. Do not place Console near sensitive equipment when operating.
- Equipment not suitable for use in the presence of flammable anesthetic mixture with air or with oxygen or nitrous oxide.
- To avoid the risk of electric shock, this equipment must only be connected to a supply mains with protective earth.
- Where the "Trapping Zone Hazard for Fingers" symbol is displayed on the console, there exists a risk of trapping or pinching fingers during operation and care must be exercised to avoid injury.
- Do not reposition or push the console from any point other than the handle designed for that purpose. A condition of overbalance or tipping may ensue.
- The AngioJet Ultra Console should not be used adjacent to or stacked with other equipment, and if adjacent or stacked use is necessary, the AngioJet Ultra Console should be observed to verify normal operation in the configuration in which it will be used.
- Portable and mobile RF communications equipment can affect MEDICAL ELECTRICAL EQUIPMENT.
- The use of accessories and cables other than those specified, with the exception of accessories and cables sold by Bayer HealthCare as replacement parts for internal components, may result in increased EMISSIONS or decreased IMMUNITY of the Ultra Console.
- MEDICAL ELECTRICAL EQUIPMENT needs special precautions regarding Electro-Magnetic Compatibility (EMC) and needs to be installed and put into service according to the EMC information provided in the tables provided in the IFU.

ADVERSE EVENTS

Potential adverse events which may be associated with use of the AngioJet Ultra Thrombectomy System are similar to those associated with other interventional procedures and include, but are not limited to:

- abrupt closure of treated vessel • acute myocardial infarction • acute renal failure • bleeding from access site • cerebrovascular accident • death • dissection • embolization • proximal or distal • hematoma • hemolysis • hemorrhage, requiring transfusion • hypotension/hypertension • infection at the access site • pain • pancreatitis • perforation • pseudoaneurysm • reactions to contrast medium • thrombosis/occlusion • total occlusion of treated vessel • vascular aneurysm • vascular spasm • vessel wall or valve damage

STERLING MR STERLING OTW

CAUTION: Federal law (USA) restricts this device to sale by or on the order of a physician. Rx only. Prior to use, please see the complete "Instructions for Use" for more information on Indications, Contraindications, Warnings, Precautions, Adverse Events, and Operator's Instructions.

INTENDED USE/INDICATIONS FOR USE

The Sterling PTA Balloon Dilatation Catheters are indicated for Percutaneous Transluminal Angioplasty in the peripheral vasculature, including iliac, femoral, popliteal, infra-popliteal, renal, and for the treatment of obstructive lesions of native or synthetic arteriovenous dialysis fistulae. These devices are also indicated for post-dilatation of balloon expandable and self-expanding stents in the peripheral vasculature. The Sterling Monorail PTA Balloon Dilatation Catheter (only) is also indicated for the carotid arteries.

CONTRAINDICATIONS

None Known.

GENERAL PRECAUTIONS

The Sterling PTA Balloon Dilatation Catheter should be used with caution for procedures involving calcified lesions or synthetic vascular grafts due to the abrasive nature of these lesions.

The Sterling PTA Balloon Dilatation Catheters are not intended for injection of contrast medium.

The Sterling™ PTA Balloon Dilatation Catheter shall only be used by physicians trained in the performance of percutaneous transluminal angioplasty.

Precautions to prevent or reduce clotting should be taken when any catheter is used:

- Consider systemic heparinization.
- Flush or rinse all products entering the vascular system with sterile isotonic saline or a similar solution prior to use.

ADVERSE EVENTS

The complications that may result from a balloon dilatation procedure include, but are not limited to:

- Allergic reaction to contrast medium • Arrhythmias • Arteriovenous fistula • Cerebrovascular accidents (specific to MR) • Death • Hematoma • Hemodynamic instability • Hemorrhage • Pseudoaneurysm • Pyrogenic reaction • Sepsis/infection • Thromboembolic episodes • Vascular thrombosis • Vessel injury, e.g. dissection, perforation, rupture • Vessel occlusion • Vessel spasm

VICTORY PERIPHERAL GUIDEWIRE

CAUTION: Federal law (USA) restricts this device to sale by or on the order of a physician. Rx only. Prior to use, please see the complete "Instructions for Use" for more information on Indications, Contraindications, Warnings, Precautions, Adverse Events, and Operator's Instructions.

INDICATIONS FOR USE:

The Victory guidewires are intended to facilitate the placement and exchange of balloon catheters or other interventional devices within the peripheral vasculature during Percutaneous Transluminal Angioplasty (PTA) or other intravascular interventional procedures.

CONTRAINDICATIONS:

The Victory guidewires are not intended for use in the coronary or cerebral vasculatures or in patients judged not acceptable for percutaneous intervention.

WARNINGS:

PRECAUTIONS:

- This device should be used only by physicians trained in percutaneous, intravascular techniques and/or procedures.
- Carefully read all instructions prior to use. Observe all warnings and precautions noted throughout these instructions. Failure to do so may compromise guidewire performance and result in complications.

ADVERSE EVENTS:

Potential adverse events which may result from use of the device include but are not limited to:

- Hematoma and other access site complications • Death • Hemorrhage (bleeding) • Reaction to contrast media • Irritation to vessel causing vessel spasm • Vessel dissection or perforation • Thrombus formation

THRUWAY .014 GUIDEWIRE

CAUTION: Federal law (USA) restricts this device to sale by or on the order of a physician. Rx only. Prior to use, please see the complete "Instructions for Use" for more information on Indications, Contraindications, Warnings, Precautions, Adverse Events, and Operator's Instructions.

INTENDED USE/ INDICATIONS FOR USE

The Thruway Guidewire facilitates placement of a catheter during diagnostic or interventional peripheral intravascular procedures including but not limited to renal intervention. The wire can be torqued to facilitate navigation through the vasculature.

CONTRAINDICATIONS

- Not intended for use in coronary arteries.
- Not intended for use in the neurovasculature.

WARNINGS/ADVERSE REACTIONS

The complications that may result from the use of a guidewire in a procedure include:

- Vessel perforation, dissection, trauma or damage • Embolism • Hematoma • Infection • Vessel spasm • Hemorrhage • Renal Failure • Myocardial Infarction • Vascular thrombosis • Stroke • Death

CholCE, CholCE PT, Luge, Mailman, PT Graphix

CholCE Magnet, CholCE PT Magnet, Luge Magnet, Mailman Magnet, PT Graphix Magnet

CAUTION: Federal law (USA) restricts this device to sale by or on the order of a physician. Rx only. Prior to use, please see the complete "Instructions for Use" for more information on Indications, Contraindications, Warnings, Precautions, Adverse Events, and Operator's Instructions.

INTENDED USE/INDICATIONS FOR USE

Boston Scientific CholCE, CholCE PT, Luge, Mailman and PT Graphix Guidewires, and Boston Scientific CholCE Magnet, CholCE PT Magnet, Luge Magnet, Mailman Magnet, and PT Graphix Magnet Guidewires with ICE Hydrophilic Coating are intended to facilitate the placement of balloon dilatation catheters or other therapeutic devices during PTCA or other intravascular interventional procedures. They are not intended for use in the cerebral vasculature.

CONTRAINDICATIONS

None known.

WARNINGS

Guidewires should be used only by physicians thoroughly trained in their intended use. Use extreme caution and careful judgment in patients for whom anticoagulation is not indicated. Severe reaction may occur in response to contrast agents that cannot be adequately premedicated. Excessive force against resistance may result in separation of the guidewire tip, damage to the catheter or vessel damage. Resulting guidewire fractures might require additional percutaneous intervention or surgery. The hydrophilic coating of these guidewires increases the possibility of vessel wall perforation compared to non-hydrophilic coatings. Care should be taken when advancing a guidewire after stent deployment. A guidewire may exit between stent struts when recrossing a stent that is not fully apposed to the vessel wall. Subsequent advancement of any device over the guidewire could cause entanglement between the guidewire and the stent.

PRECAUTIONS

Carefully check and match therapeutic device compatibility to the wire prior to use. Sharp insertion tools may compromise the integrity of the polymer coating. To avoid guidewire damage and possible shearing of plastic, do not withdraw or manipulate the wire through a metal needle cannula. Excessive tightening of the torque device onto the wire may result in abrasion of the coating on the wire.

Boston Scientific guidewires (DFU 90976970) are designed to be compatible exclusively with the AddWire™ Extension Wire for interventional device exchange. Do not use another extension or exchange system. Carefully check and match the compatibility of the guidewire diameter with the interventional device prior to use. These guidewires should only be used in devices having an inner lumen diameter greater than 0.015 in (0.39 mm).

Boston Scientific Magnet guidewires (DFU 90976935) should only be used in devices having an inner lumen diameter greater than 0.015 in (0.39 mm). **NOTE THAT THE ACTUAL DIAMETER MAY BE UP TO 0.015 IN (0.39MM) IN THE MAGNET EXCHANGE SEGMENT OF 182 CM guidewires.** Compatibility of the therapeutic device with the 182 cm guidewire and The MAGNET Exchange Device should be verified prior to use to ensure holding force adequate to keep wire from moving during exchange.

ADVERSE EVENTS

Potential adverse events which may result from the use of the device include but are not limited to:

- Allergic reaction to contrast media • Embolism • Hemorrhage or hematoma • Infection, local infection, systemic infection • Pain at the access site • Pseudoaneurysm • Vascular thrombus • Vessel spasm • Vessel trauma (dissection, perforation, rupture or injury)

In addition, when used for PTCA:

- Abrupt closure • Angina or unstable angina • Arrhythmias • Cardiac tamponade/pericardial effusion • Contrast induced renal insufficiency or renal failure • Death • Myocardial infarction or ischemia • Stroke/ cerebral vascular accident (CVA)/transient ischemic attack (TIA)

Some of the above potential adverse events may require additional surgical intervention.

All Trademarks are the property of their respective owners. © 2015 Boston Scientific Corporation or its affiliates. All rights reserved. PI-347609-AA NOV2015