

Chemoembolization Via Intrahepatic Collateral Arteries

BY DANIEL Y. SZE, MD, PhD

CASE SUMMARY

A 53-year-old man with chronic hepatitis B presented with elevated transaminases while taking herbal supplements, but was otherwise asymptomatic, Child-Pugh A5, and ECOG performance status 0. Workup included nonreactive hepatitis B surface antigen, negative hepatitis C antibody, negative colonoscopy except for small hemorrhoids, negative antinuclear antibody, negative alpha-1 antitrypsin, and ceruloplasmin tests. He transferred his care to a second hospital and underwent hepatic ultrasonography, revealing a 16-cm mass, confirmed on multiphase CT scan to be a hepatocellular carcinoma (Figure 1). He then transferred his care to a tertiary care hospital, where he underwent a trisegmentectomy.

One year after “curative” resection, follow-up imaging revealed a hypervascular lesion in segment 3 measuring 2.1 cm, indicative of recurrence (Figure 2). A second possible lesion in the fissure of the ligamentum venosum was measured at 0.8 cm and was felt to be inaccessible for ablation (Figure 2). The patient’s liver function remained Child-Pugh A5, but his performance

status had deteriorated mildly to status 1 after surgery. He was referred for hepatic angiography and chemoembolization.

IMAGING FINDINGS

Initial hepatic angiography revealed postsurgical distortion of the arterial anatomy, with proximal occlusion of the segment 2 and segment 3 arteries with distal reconstitution from intrahepatic collateral vessels (Figure 3). Parenchymal phase imaging confirmed two hypervascular lesions (Figure 3).

Although the collateral vessels appeared too small in diameter and too tortuous to accommodate standard




Figure 1. Coronal reformat of arterial-phase CT shows replacement of the right lobe with a large hepatocellular carcinoma, extending into segments 1 and 4.




Figure 2. Arterial-phase CT image of the remnant liver 1 year after resection shows a new 2.1-cm lesion in segment 3 (arrow). Washout and pseudocapsule confirmed recurrent hepatocellular carcinoma (A). A second small nodule of hypervascularity is seen in the fissure of the ligamentum venosum adjacent to the left portal vein (arrow) but is too small to characterize further (B).




Figure 3. Left hepatic arteriogram revealed occlusion of the proximal segment 2 and segment 3 branches, with reconstitution of distal branches via intrahepatic collateral vessels at the cut surface (A). Parenchymal phase image from the same arteriogram confirmed two hypervascular lesions (B).




Figure 4. Selective arteriogram of the most cranial segment 2 collateral branch confirmed supply to the smaller lesion. A 2.8-Fr SeQuire® catheter was wedged into this submillimeter vessel, and harder injection showed additional collateralization reconstituting distal segment 2 vessels (A). Selective arteriogram of the largest but most tortuous collateral vessel reached the reconstituted segment 3 vessel, supplying the larger tumor (B).




Figure 5. After administration of 100–300-µm doxorubicin-eluting microspheres at both sites, the final arteriogram showed resolution of tumor blushes and retained patency of the collateral network, segment 2 artery, and proximal segment 3 artery (A). Unenhanced cone-beam CT showed retention of iodinated contrast medium in the tumors, as well as in some peripheral arteries (B).

microcatheters, treatment was attempted using a Fathom 0.014-inch guidewire (Boston Scientific Corporation) and a 2.8-Fr SeQuire® microcatheter (Guerbet). (At the time, smaller-diameter SeQuire® microcatheters had not yet been released.)

The submillimeter branch supplying the falciform cleft was successfully selected, and the lesion was treated with doxorubicin-loaded LC beads 100–300 µm (BTG) until stasis of the tumor-feeding branch (Figure 4) was achieved. Likewise, the tortuous intrahepatic collateral vessel to segment 3 was selected and the segment 3 lesion was also treated with drug-eluting LC beads (Figure 4). Completion angiography confirmed disappearance of tumor blushes but maintenance of flow in nontarget vessels (Figure 5), and unenhanced cone-beam CT confirmed retention of contrast medium in the treated lesions (Figure 5).

Follow-up MRI at 3 months revealed complete response of the ligamentum venosum lesion and partial response of the segment 3 lesion. Residual disease was treated with percutaneous microwave ablation.

DISCUSSION

Microcatheter design must balance the demands of: (1) adequate lumen to accommodate viscous or particle-containing injected fluids; (2) flexibility to navigate tortuosities without causing vessel injury; and (3) longitudinal stiffness to allow pushability. Additional features may include shaped tips, torqueability, and characteristics that help reduce reflux.

CONCLUSION

Selectivity of catheterization and treatment has a substantial impact on outcomes of transarterial chemoembolization. Navigation of small, tortuous vessels is possible using the SeQuire® microcatheter, even using the large 2.8-Fr version. Although this case cannot demonstrate the reflux reduction features of the SeQuire® microcatheter, it demonstrates the flexibility and pushability that allowed selection of small, tortuous vessels. Future studies using

radiopaque microspheres may be performed to test the controlled reflux feature. ■

Recommended Reading

1. Miyayama S, Mitsui T, Zen Y, et al. Histopathological findings after ultraslective transcatheter arterial chemoembolization for hepatocellular carcinoma. *Hepatol Res.* 2009;39:374–381.
2. Mokin M, Waqas M, Setlur Nagesh SV, et al. Assessment of distal access catheter performance during neuroendovascular procedures: measuring force in three-dimensional patient specific phantoms. *J Neurointerv Surg.* 2019;11:619–622.
3. Ogata N, Goto K, Uda K. An evaluation of the physical properties of current microcatheters and guidewires. The development of the “catheter-glide approach” in response to weaknesses of current materials. *Interv Neuroradiol.* 1997;3:65–80.
4. Zoarski GH, Mathis JM, Hebel JR. Performance characteristics of microcatheter systems in a standardized tortuous pathway. *AJNR Am J Neuroradiol.* 1998;19:1571–1576.

Daniel Y. Sze, MD, PhD

Professor of Vascular and Interventional Radiology
Stanford University
Stanford, California