Body Dysmorphic Disorder Screening Tools for the Dermatologist: A Systematic Review

BY MELISSA DANESH, BS, KOUROSH BEROUKHIM, BS, CATHERINE NGUYEN BS, ETHAN LEVIN MD, AND JOHN KOO MD

ody Dysmorphic Disorder is a psychiatric condition defined by distressing and/or impairing preoccupation with a non-existent or slight defect in appearance. The most recent diagnostic criteria have added the additional criterion of repetitive behaviors or mental acts in response to preoccupations with perceived defects of flaw in physical appearance. The most common areas of concern are the skin (e.g., minimal acne), hair (e.g., thinning), and the nose.²⁻⁴

Screening questionnaires for BDD have identified a substantial percentage of patients with BDD who present to dermatology clinics. In a survey of 268 patients seeking dermatological treatment, a total of 11.9 percent (95 percent Confidence Interval [CI], 8.0-15.8 percent) of patients screened positive for BDD. Rates were similar in a community general dermatology setting (14.4 percent) and a university cosmetic surgery setting (10 percent), making dermatologists the physicians most often seen by patients with BDD.5 Despite its prevalence in the dermatology setting, the disease remains underdiagnosed. 6-10 Hence, physicians may attempt to fix the perceived flaw, but these patients are often dissatisfied and may sue or become violent toward the treating physician. 11-14 Screening for BDD may thus be warranted before patients undergo cosmetic procedures.

The issue is further underscored by the morbidity and mortality associated with BDD: patients often have concomitant depression and anxiety, withdrawal from

relationships and social activities, suicide attempts, and completed suicide.^{2,11,13,15,16} The morbidity of this disease has been illustrated within the dermatology practice as well after a study showed sixteen dermatology patients who committed suicide, most of whom had either acne or BDD. 16

A systematic review of screening tools for BDD was recently done, however it broadly discussed screening tools used in a variety of patient populations (cosmetic surgery, dermatology, rhinoplasty, orthognathic surgery, and cosmetic dental) and only provided actual questions from one of many screening tools discussed. The aim of this review is to comprehensively review screening tools for BDD that have been used or validated in the dermatological setting. We will also show some of the actual screening tools and a description on how to use each of them in order to facilitate their utilization in the dermatology clinic.

PRACTICAL POINTER

Currently, many of patients with Body Dysmorphic Disorder (BDD) are being treated suboptimally without psychiatric care. Early screening and diagnosis can aid in the difficult management involved in caring for these patients, and more importantly, may reduce the severe morbidity and mortality associated with BDD.

OUR REVIEW

An electronic PubMed search was conducted to identify all screening tools for BDD in the dermatology patient population. After eliminating duplicate studies and non-English written studies, the search result was evaluated by the first two authors. Irrelevant items were excluded after reviewing the titles or abstracts of the all identified articles. Full text articles were then evaluated to determine if they met eligibility criteria. Inclusion criteria included 1) BDD was defined according to criteria defined in the DSM-IV or DSM-V; 2) The paper utilized a screening tool for diagnosing BDD and 3) The study investi-

gated a population seeking dermatological treatment. Our electronic search yielded 151 articles, of which 48 full text articles were retrieved. We identified six studies (Table 1) that assessed the presence of BDD according to

its definition in the DSM-IV and that were use in the dermatology setting. Within these six studies, we identified

three different screening tools.

SCREENING INSTRUMENTS

Body Dysmorphic Disorder Questionnaire-Dermatology Version

The Body Dysmorphic Disorder Questionnaire-Dermatology Version (Appendix 1) is a modified version of the Body Dysmorphic Disorder Questionnaire (BDDQ), which has been validated for use in the psychiatric setting (sensitivity 100 percent, specificity 89 percent). The original BDDQ is based on the definition of BDD provided in the Diagnostic and Statistical Manual of Mental Disorders-IV (DSM-IV) and consists of four sets of "yes/no" questions.¹⁷ The Body Dysmorphic Disorder Questionnaire-Dermatology Version (BDDQ-DV; Appendix 1) was developed by Dufresne et al. and Phillips et al., involving the following modifications: substitution of a Likert scale from 1-5 to indicate a range of severity, rather than "yes/no" responses, and the removal of a question on the effects of the patient's concern on family or friends.^{5, 14} To screen positive for BDD, patients must report the presence of preoccupation as well as at least moderate (score of 3 or higher) distress or impairment in functioning.

	ry concerned about the appearance of specially unattractive?	Y	N					
If no, thank	If no, thank you for your time and attention. You are finished with this questionnaire.							
	ese concerns preoccupy you? That is,	V						
naro to stop	thinking about?		N					
What are these concerns? What specifically bothers you about the appearance of these body parts?								
What effect has your preoccupation with your appearance had on your life?								
Has your defect often caused you a lot of distress, torment or pain? How much? (circle best answer)								
1	2	3	4	5				
No		Moderate and disturbing but still	Severe, and very	Extreme, and				
distress	Mild, and not too disturbing	manageable	disturbing	disabling				
Has your defect caused you impairment in social, occupational or other important areas of functioning? How much? (circle best answer)								
1	2	3	4	5				
No	Mild interference but overall per-	Moderate, definite interference, but	Severe, causes substantial	Extreme,				
limitation	formance not impaired	still manageable	impairment	incapacitating				
Has your de	efect often significantly interfered with y	our social life?	Υ Υ	N				
If yes, how?	}							
Has your defect often significantly interfered with your school work, your job, or your								
ability to function in your role?			Υ .	N				
Are there things you avoid because of your defect?			Y	N				

Appendix 1. Body Dysmorphic Disorder Questionnaire-Dermatology Version (BDDQ-DV)

The BDDQ-DV has been used in four studies to assess BDD in a dermatological setting, one in patients seeking treatment for acne vulgaris, 18 and three involving patients who presented to cosmetic or general dermatology practices.5,14,19 This BDDQ-DV was reported to have comparable sensitivity (100 percent) and slightly improved specificity (94.7 percent) compared to the original questionnaire validated for the psychiatric setting. 14 Additionally, in the study by Bowe et al. the majority of patients who screened positive for BDD provided detailed descriptions of their distress and impairment, providing further validation of this tool. Two of the studies also used the BDDQ-DV to assess for BDD in the general dermatology clinic.^{5,19}

Dysmorphic Concern Questionnaire

The Dysmorphic Concern Questionnaire (DCQ; Appendix 2) is a practical, seven-item questionnaire, which was developed and validated by Oosthuizen for the psychiatric setting. (Appendix 2) It is based on the General Health Questionnaire (GHQ), an instrument that was devised to quantify the risk of developing psychiatric disorders, which measures common mental health problems including depression, somatic symptoms, and social withdrawal.²⁰ The DCQ is focused on BDD, and asks about patient concern with physical appearance and past attempts to deal with the issue. Each item is answered on a four-point scale (answers ranging from 0-3 points). The DCQ was found to have good internal consistency (Chronbach's alpha: .88) and validity as demonstrated by strong correlations with distress, work, and social impairment.²¹

Stangier et al22 validated the DCQ as a screening instrument in a dermatological sample of 156 female outpatients who presented for both cosmetic and non-cosmetic treatment. The study found an internal consistency of Cronbach's alpha: .85 (slightly decreased from it's use in the psychiatric setting), and set a cut-off value at ≥14 to achieve maximum sensitivity (72 percent) and specificity (90.7 percent).

Body Dysmorphic Symptom Scale

The Body Dysmorphic Symptoms Scale (BDSS) is a 10-item self-reported questionnaire, which asks about patient concerns with appearance and related behaviors (i.e., looking

in the mirror repeatedly), and the impact of these perceived defects on patients' social life and relationships.

A patient's score is obtained by the sum of each positive question. There is no accepted cut-off point for the BDSS in order to screen positive for BDD.

Dysmorphic Concerns Questionnaire

These questions are designed to screen for people with certain concerns that are: (i) often difficult or embarrassing to talk about with their doctor/family/friends; and (ii) often difficult to find the right help for.

Please read the following questions carefully and answer them by circling the answer which you think is most appropriate for your specific situation:

Have you ever:				
Been very concerned about some aspect of your physical appearance	Not at all	Same as most people	More than most people	Much more than most people
 Considered yourself misformed or misshapen in some way (e.g. nose/hair/skin/sexual organs/ overall body build) 	Not at all	Same as most people	More than most people	Much more than most people
 Considered your body to be malfunctional in some way (e.g. excessive body odour, flatulence, sweating) 	Not at all	Same as most people	More than most people	Much more than most people
 Consulted or felt you needed to consult a plastic surgeon/dermatologist/physician about these concerns 	Not at all	Same as most people	More than most people	Much more than most people
Been told by others/doctor that you are normal in spite of you strongly believing that something is wrong with your appearance or bodily functioning	Not at all	Same as most people	More than most people	Much more than most people
 Spent a lot of time worrying about a defect in your appearance/bodily functioning 	Not at all	Same as most people	More than most people	Much more than most people
7. Spent a lot of time covering up defects in your appearance/bodily functioning	Not at all	Same as most people	More than most people	Much more than most people

Appendix 2. Dysmorphic Concern Questionnaire (DCQ)

Kaymak et al.²³ used the BDSS in 107 Turkish university students diagnosed with skin disease at an outpatient dermatology clinic. The study used a score of ≥4 as the score highly favoring a diagnosis of BDD. The study did not use any additional tools or a structured clinical inter-

TABLE 1								
BDD Screening Tools Applied in a Dermatology Setting								
Questionnaire Used	Population	No. of Patients	Study					
BDDQ-DV	Outpatient cosmetic and laser surgery clinic	46	Dufresne et all, (REF) 2001					
BDDQ-DV	General dermatology and dermatological cosmetic surgery	268	Phillips et al, (REF) 2000					
DCQ	Female dermatological outpatients	156	Stangier et al, 2003					
BDDQ-DV	Acne Vulgaris	128	Bowe et al, 2007					
BDSS	University student dermatological outpatients	107	Kaymak et al, 2009					
BDDQ-DV	Cosmetic and general dermatology clinic	400	Dogruk et al, 2014					

view to confirm the diagnosis of BDD in patients who screened positive using the BDSS. As such, the study did not attempt to validate the BDSS to screen BDD.

DISCUSSION

Three screening tools have been utilized to diagnose BDD (as defined by DSM-IV or DSM-V) in the dermatological setting: BDDQ-DV, DCQ, and BDDS. The BDDQ-DV and DCQ have also been validated in this setting. All three tools have a limited number of items, making them practical for the outpatient dermatology setting. However, the BDDQ-DV has been used most commonly in dermatology, perhaps making it the most reliable of the three tools.

BDD does not affect any specific "type" of patient, and thus assessing those at risk should include screening tools to create a more standardized method for diagnosis. Despite an onset in adolescence,²⁴ establishing a diagnosis may take 10-15 years. This may be, at least in part, due to difficulty recognizing high-risk patients based on specific treatments they are seeking or demographic information. For instance, while one may assume patients seeking cosmetic treatments have a significantly higher rate of BDD, recent data⁵ has indicated that the rate of BDD does not significantly differ between patients who present to cosmetic (10.0 percent) and general dermatology (14.4 percent) clinics. Additionally, BDD may be as common in men as in women, eliminating gender as a way to identify at risk patients. 12 Hence, the availability of a practical, accurate tool to screen for and diagnose BDD, which has proven effective in the dermatological setting, is important in identifying and diagnosing these patients.

There remains great need for further research to advance our management and reporting of dermatological patients with BDD, given the demonstrated prevalence of these patients in the dermatology setting. Currently, many of these patients are being treated suboptimally without psychiatric care. The early screening and diagnosis can aid in the difficult management involved in caring for these patients, and more importantly, may reduce the severe morbidity and mortality associated with BDD.

The authors have no conflict of interest to declare.

There have been no prior presentations of this material.

Melissa Danesh is a UCSF medical student currently doing research at the Psoriasis, Phototherapy & Skin Treatment Clinic at UCSF under Dr. John Koo, who is double-boarded in both dermatology and psychiatry.



Kourosh Beroukhim, Department of Dermatology, University of California, San Francisco, San Francisco.

Catherine Nguyen, Department of Dermatology, University of California, San Francisco, San Francisco.

Ethan Levin, MD, Department of Dermatology, University of California, San Francisco, San Francisco.

John Koo, MD, Director of the Psoriasis, Phototherapy, and Skin Treatment Clinic, Department of Dermatology, University of California, San Francisco, is double-boarded in both dermatology and psychiatry.



- 2. Phillips KA, McElroy SL, Keck PE, Pope HG, Hudson JI: Body dysmorphic disorder: 30 cases of imagined ugliness. American journal of Psychiatry 1993, 150:302-.
- 3. Hollander E, Cohen L, Simeon D: Body dysmorphic disorder. Psychiatric Annals 1993, 23:359-64
- 4. Veale D, Boocock A, Gournay K, Dryden W, Shah F, Willson R, Walburn J: Body dysmorphic disorder. A survey of fifty cases. The British Journal of Psychiatry 1996, 169:196-201.
- 5. Phillips KA, Dufresne Jr RG, Wilkel CS, Vittorio CC: Rate of body dysmorphic disorder in dermatology patients. Journal of the American Academy of Dermatology 2000, 42:436-41.
- 6. Cotterill J: Dermatological non disease: a common and potentially fatal disturbance of cutaneous body image. British Journal of Dermatology 1981, 104:611-9.
- 7. Bishop Jr ER: Monosymptomatic hypochondriacal syndromes in dermatology. Journal of the American Academy of Dermatology 1983, 9:152-8.
- 8. Koblenzer C: The broken mirror: dysmorphic syndrome in the dermatologist's practice. Fitz J Clin Dermatol 1994,
- 9. Zaidens SH: Dermatologic hypochondriasis: a form of schizophrenia. Psychosomatic medicine 1950, 12:250-3. 10. Hanes KR: Body dysmorphic disorder: an underestimated entity? Australasian Journal of Dermatology 1995, 36:227-8
- 11. Cotterill JA: Body dysmorphic disorder. Dermatologic clinics 1996, 14:457-63.
- 12. Phillips KA, Diaz SF: Gender differences in body dysmorphic disorder. The Journal of nervous and mental disease
- 13. Phillips KA: The broken mirror: understanding and treating body dysmorphic disorder: Oxford University Press,
- 14. Dufresne RG, Phillips KA, Vittorio CC, Wilkel CS: A screening questionnaire for body dysmorphic disorder in a cosmetic dermatologic surgery practice. Dermatologic surgery: official publication for American Society for Dermatologic Surgery et al. 2001, 27:457-62.
- 15. Phillips KA: Body dysmorphic disorder: the distress of imagined ugliness. Am J Psychiatry 1991, 148:1138-49. 16. Cotterill J, Cunliffe W: Suicide in dermatological patients. British Journal of Dermatology 1997, 137:246-50.
- 17. Phillips K, Atala K, Pope H: Diagnostic instruments for body dysmorphic disorder. 1995 Annual Meeting New Research Program and Abstracts Washington, DC, American Psychiatric Association, 1995.
- 18. Bowe WP, Leyden JJ, Crerand CE, Sarwer DB, Margolis DJ: Body dysmorphic disorder symptoms among patients with acne vulgaris. Journal of the American Academy of Dermatology 2007, 57:222-30.
- 19. Dogruk Kacar S, Ozuguz P, Bagcioglu E, Coskun KS, Uzel Tas H, Polat S, Karaca S: The frequency of body dysmorphic disorder in dermatology and cosmetic dermatology clinics: a study from Turkey. Clinical and experimental dermatology 2014, 39:433-8.
- 20. Goldberg DP: The detection of psychiatric illness by questionnaire: A technique for the identification and assessment of non-psychotic psychiatric illness. 1972.
- 21. Oosthuizen P, Lambert T, Castle DJ: Dysmorphic concern: prevalence and associations with clinical variables. Australian and New Zealand Journal of Psychiatry 1998, 32:129-32.
- 22. Stangier U, Janich C, Adam-Schwebe S, Berger P, Wolter M: Screening for body dysmorphic disorder in dermatological outpatients. Dermatology and Psychosomatics/Dermatologie und Psychosomatik 2003, 4:66-71.
- 23. Kaymak Y, Taner E, im ek I: Body dysmorphic disorder in university students with skin diseases compared with healthy controls. Acta dermato-venereologica 2009, 89:281-4.
- 24. Phillips KA, Menard W, Fay C, Weisberg R: Demographic Characteristics, Phenomenology, Comorbidity, and Family History in 200 Individuals With Body Dysmorphic Disorder. Psychosomatics 2005, 46:317-25.







