




FIGURE 1

CATCHING FAMILIAL EXUDATIVE VITREORETINOPATHY EARLY



Know the first signs of this inherited retinal disease to initiate prompt treatment and monitoring.

BY VAIDEHI SATHAVE, MS; MANISH NAGPAL, MS, FRCS, FASRS; AND NAVNEET MEHROTRA, DNB, FRF

A 13-year-old boy was brought to our clinic for a routine visit. His birth history was normal, and there was no significant family history. On examination, his VA was 20/20 OU. His IOP was 20 mm Hg OD and 19 mm Hg OS. The anterior segment examination was within normal limits. Dilated fundus examination revealed avascular ischemic areas in the periphery of each eye, with a normal posterior pole (Figure 1).

Fluorescein angiography showed 360° capillary nonperfusion in the periphery with leakage secondary to neovascular proliferations in the temporal periphery of each eye (Figure 2).

A diagnosis of stage 1 familial exudative vitreoretinopathy (FEVR) was made, and the patient was treated with laser photocoagulation. At the 1-month follow-up, each eye

showed a well-lasered retinal periphery (Figure 3). The patient will be evaluated again 4 to 6 months post-treatment.

STAGING

FEVR is a hereditary vitreoretinal disorder in which there is abnormal retinal angiogenesis, resulting in incomplete peripheral retinal vascularization and retinal ischemia. A clinical classification describes five stages of FEVR¹:

- Stage 1:** avascular periphery or anomalous intraretinal vascularization
- Stage 2:** avascular retinal periphery with extraretinal vascularization

(Continued on page 49)

(Continued from page 50)

FIGURE 2





FIGURE 3



Stage 3: extramacular retinal detachment (RD)

Stage 4: macula-involving, subtotal RD

Stage 5: total RD

Stage 1 FEVR often goes undiagnosed due to the lack of symptoms. FEVR usually has a progressive course during childhood and may become stable after 20 years of age; however, late progression with vision-threatening complications, such as vitreous hemorrhage and RD, can occur at any age.

MANAGEMENT

Treatment protocols include observation, laser photocoagulation, and surgery (ie, vitrectomy or scleral buckling), depending on the disease stage. Long-term monitoring is required. Examination of the family members is also critical so that early intervention can be planned before complications develop. ■

1. Kashani AH, Learned D, Nudleman E, Drenser KA, Capone A, Trese MT. High prevalence of peripheral retinal vascular anomalies in family members of patients with familial exudative vitreoretinopathy. *Ophthalmology*. 2014;121(1):262-268.

VAIDEHI SATHAYE, MS

- Retina and Vitreous Fellow, The Retina Foundation, Ahmedabad, India
- vaidehisathaye@gmail.com
- Financial disclosure: None

MANISH NAGPAL, MS, FRCS, FASRS | SECTION EDITOR

- Senior Consultant, Retina and Vitreous Services, The Retina Foundation, Ahmedabad, India
- drmanishnagpal@yahoo.com
- Financial disclosure: Consultant (Nidek)

NAVNEET MEHROTRA, DNB, FRF

- Vitreoretinal and Uveitis Consultant, The Retina Foundation, Ahmedabad, India
- Financial disclosure: None

**If you have images you would like to share, email
Dr. Nagpal at drmanishnagpal@yahoo.com.**

Note: Photos should be 400 dpi or higher and at least 10 inches wide.